

The present study investigates the actual problem of determining the psychological characteristics and content of fears among modern Ukrainian students-practical psychologists. The importance of studying this problem today lies in the fact that the vast majority of people in Ukrainian society is under stress caused by events occurring in the country. The processes of vital self-determination of youth that revolve around the problem of value and affordability is also powerful factor of stress. We assume that the social context of fears, and the presence of gender- professional features of the fears among students-practical psychologists.

Nosova, Eugenia

Shupyk National Medical Academy of Postgraduate Education, Kyiv

Features of cognitive-emotional deficits in patients with schizophrenia with manic syndrome

Recently, the attention of psychologists and psychiatrists focused on attempts to quantify mental disorders. This includes the MSCEIT questionnaire, which determines the degree of development of emotional intelligence. As has been established, social adaptation of patients with schizophrenia is disrupted to a greater extent due to dysfunction in the sphere of recognition and processing of emotions. Intelligence, for a long time retains its performance. Additionally pronounced decrease in all branches of the questionnaire in schizophrenia can occur at the expense of the violations in the emotional sphere (apathetic changes etc.), and thought disorder in the form of violations of generalization and distraction. Characteristic of schizophrenia schisis affects the emotional side of the psyche. In this regard, decided classical technique MSCEIT to complement the Wechsler test, the methodology of the icons and the method of interpretation of Proverbs and metaphors in modification B. V. Zeigarnik, the purpose of a comprehensive assessment of psycho-pathological disorders. And also to identify potential resource areas in patients with schizophrenia required for the subsequent rehabilitation and adaptation. For study will be selected group of patients with schizophrenia coupled with manic syndrome and healthy subjects for control. The results will be presented to attention of participants of conference.

Oprya, Yevgen

Odessa National Medical University

Many persons are suffering from serious chronic mental illness, such as schizophrenia and bipolar disorder, which has poor response to pharmacotherapy, bad prognosis and leads to disability. Retired mental patients and their families experience severe distress due to patient's disability. Purpose of work is how to best provide complex treatment-prophylaxis help to these patients by using standards of palliative care to support best quality of their life by decrease emotional stress and improve of social functioning. It is planned to assess level of stress (anxiety) by using HAM-A scale and establish correlation between level of stress and EEG changes (index of activation (the ratio of beta and alpha indexes).

Prutko, Kateryna

Kharkiv National Medical University

Oneirofrenia

My project is researching state called Oneirofrenia, it was discovered by L.J.Meduna. Oneirofrenia is a syndrome that in the overwhelming majority of

cases is connecting with schizophrenia pathology, sometimes it can be single pathology. The basic symptom is a disturbance of apperception. The sense modalities mostly affected, in the order of frequency are vision, various proprioceptions and interoception, including body image, smell, hearing. Prognosis is usually good.

Medical case: The patient was in acute state, had a premorbid. Had a persecutory delusion, visual hallucination, she sad about that state "everything was different I was like Alice in wonderland», «I was so scared I can't recognize my husband, children. The colors was different, I can't trust my eyes. "She said that she spoke with her mother and she asked her to take her children to the church, don't explain for what. Woman took her children to the church, and start to screaming there. When the husband took her home she has fallen asleep. When she got up she didn't remember some things that happening yesterday. Also I have her diary with the picture that she draw at this condition. I'm pretty sure that if this diagnoses will be better known in psychiatry practice, for many people we can keep quality of life.

Rusanov, Vladimir

Bogomolets National Medical University, Kyiv

ADHD and role models for children in modern society

In the field of my scientific interests are humanistic and social psychology and psychotherapy. I am interested in classical works by Maslow, Shostrom and Fromm but just watching the modern trends. Also, issues of pedagogy and education of children with hyperkinetic disorders (F90). And first of all, as a resident of Donetsk forced to leave own city, I'm interested in the issue of Posttraumatic stress disorder during the war and how to educate a new generation of Ukrainians, on well-known to all Europeans principles of "liberty, equality, fraternity".

Ryabukhin, Konstantin

Odessa National Medical University

Disharmonious features of gender status in women as predictors of alcohol dependence

A scientific analysis of the factors that precede the development of alcohol dependence in women found the main predictors of female alcohol addiction:

Biological: somatotype asthenic (29.18%), stenoplastyc (24.91%), piknic (14.59%) and mezoplastyc (13.88%), the rate of development - somatosexual (74.02 %) and psychosexual (76.16%); asynchrony of sexual development (55.17%) of the surveyed women with alcohol dependence;

Psychological - the predominance of accentuated features of character and gender psychotype of women (hypermasculine, hyperfeminine or infantile).

The program of complex medical-psychological correction was built considering the investigated results.

Rzaeva, Roksana

Military Medical Center of the Northern Region (Kharkiv)

The case of dissociative variant of post-traumatic stress disorder

This clinical case shows how close post-traumatic stress disorder is to psychotic symptoms. The clinical picture of the acute period of dissociative depersonalization in the absence of data history barely distinguishable from endogenous psychosis clinic. The presence of such symptoms in patients with