



Wiadomości Lekarskie

Official journal of the Polish Medical Association



Memory of
dr Władysław
Biegański

VOLUME LXXIV, ISSUE 9 PART 1, SEPTEMBER 2021

Since 1928



ALUNA Publishing House

Wiadomości Lekarskie is abstracted and indexed in: PUBMED/MEDLINE, SCOPUS, EMBASE, INDEX COPERNICUS, POLISH MINISTRY OF EDUCATION AND SCIENCE, POLISH MEDICAL BIBLIOGRAPHY

Copyright: © ALUNA Publishing House.

Articles published on-line and available in open access are published under Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.

Wiadomości Lekarskie monthly journal

You can order the subscription for the journal from Wydawnictwo Aluna by:

prenumerata@wydawnictwo-aluna.pl
Wydawnictwo Aluna
Z.M. Przesmyckiego 29
05-510 Konstancin-Jeziorna
Poland

Place a written order first.

If you need, ask for an invoice.

Payment should be done to the following account of the Publisher:

account number for Polish customers (PLN):

82 1940 1076 3010 7407 0000 0000

Credit Agricole Bank Polska S. A., SWIFT: AGRIPLP

account number for foreign customers (EURO):

57 2490 0005 0000 4600 7604 3035

Alior Bank S. A.: SWIFT: ALBPPLPW

Subscription of twelve consecutive issues (1-12):

Customers in Poland: 360 PLN/year

Customers from other countries: 320 EURO/year



Wiadomości Lekarskie

Editor in-Chief:

Prof. Władysław Pierzchała

Deputy Editor in-Chief:

Prof. Aleksander Sieroń

Statistical Editor:

Dr Lesia Rudenko

Managing Editor:

Agnieszka Rosa – amarosa@wp.pl

International Editorial Office:

Lesia Rudenko (editor) – l.rudenko@wydawnictwo-aluna.pl

Nina Radchenko (editor's assistant)

– n.radchenko@wydawnictwo-aluna.pl

Polish Medical Association (Polskie Towarzystwo Lekarskie):

Prof. Waldemar Kostewicz – President PTL

Prof. Jerzy Woy-Wojciechowski – Honorary President PTL

International Editorial Board – in-Chief:

Marek Rudnicki

Chicago, USA

International Editorial Board – Members:

Kris Bankiewicz	San Francisco, USA	George Krol	New York, USA
Christopher Bara	Hannover, Germany	Krzysztof Łabuzek	Katowice, Poland
Krzysztof Bielecki	Warsaw, Poland	Henryk Majchrzak	Katowice, Poland
Zana Bumbuliene	Vilnius, Lithuania	Ewa Małecka-Tendera	Katowice, Poland
Ryszarda Chazan	Warsaw, Poland	Stella Nowicki	Memphis, USA
Stanislav Czudek	Ostrava, Czech Republic	Alfred Patyk	Gottingen, Germany
Jacek Dubiel	Cracow, Poland	Palmira Petrova	Yakutsk, Russia
Zbigniew Gasior	Katowice, Poland	Krystyna Pierzchała	Katowice, Poland
Andrzej Gładysz	Wroclaw, Poland	Tadeusz Płusa	Warsaw, Poland
Nataliya Gutorova	Kharkiv, Ukraine	Waldemar Priebe	Houston, USA
Marek Hartleb	Katowice, Poland	Maria Siemionow	Chicago, USA
Roman Jaeschke	Hamilton, Canada	Vladyslav Smiiianov	Sumy, Ukraine
Andrzej Jakubowiak	Chicago, USA	Tomasz Szczepański	Katowice, Poland
Oleksandr Katrushov	Poltava, Ukraine	Andrzej Witek	Katowice, Poland
Peter Konturek	Saalfeld, Germany	Zbigniew Wszolek	Jacksonville, USA
Jerzy Korewicki	Warsaw, Poland	Vyacheslav Zhdan	Poltava, Ukraine
Jan Kotarski	Lublin, Poland	Jan Zejda	Katowice, Poland

Distribution and Subscriptions:

Bartosz Guterman prenumerata@wydawnictwo-aluna.pl

Graphic design / production:

Grzegorz Sztank

www.red-studio.eu

Publisher:

ALUNA Publishing House

ul. Przesmyckiego 29,

05-510 Konstancin – Jeziorna

www.wydawnictwo-aluna.pl

www.wiadomoscilekarskie.pl

www.wiadlek.pl

FOR AUTHORS

- The monthly "Wiadomości Lekarskie" Journal is the official journal of the Polish Medical Association. Original studies, review papers as well as case reports are published.
- The publication of the manuscript in "Wiadomości Lekarskie" is paid. The cost of publishing the manuscript is PLN 1,000 plus 23% VAT (for foreign authors 250 Euro). If the first author of the manuscript is a member of the Editorial Board, we do not charge a fee for printing the manuscript, and if she or he is the next co-author – the fee is PLN 500 plus 23% VAT. The publisher issues invoices. The fee should be paid after receiving positive reviews, and before publishing the manuscript. Membership of the Polish Medical Association with documented paid membership fees for the last 3 years is also the exempt from publication fee.
- Only papers in English are accepted for publication. The editors can help in finding the right person for translation or proofreading.
- Papers should be sent to the editor via the editorial panel (Editorial System), available on the journal's website at <https://www.wiadlek.pl>. In order to submit an article, free registration in the system is necessary. After registration, the author should follow the instructions on the computer screen.
- All editorial work is under control and using the editorial panel. This applies in particular to sending manuscripts, correspondence between the editor and author and the review process. In special cases, the editor may agree to contact outside the panel, especially in case of technical problems.
- Acceptable formats for individual elements of the article are as follows:
 - Content of the article – doc, docx, rtf, odt.
 - Tables – doc, docx, rtf, odt
 - Figures – JPG, GIF, TIF, PNG with a resolution of at least 300 dpi
 - Captions for figures and tables.
 These elements are sent to the editor separately using the editorial panel. References and article metadata such as titles, keywords, abstracts etc. are supplemented by the author manually in the editorial panel in appropriate places.
- The volume of original papers – including figures and references – must not exceed 21,600 characters (12 pages of typescript), and review papers – up to 28,800 characters (16 pages).
- The original manuscript should have the following structure: Introduction, Aims, Material and methods, Results, Discussion and Conclusions which cannot be a summary of the manuscript.
- When using abbreviations, it is necessary to provide the full wording at the first time they are used.
- In experimental manuscripts in which studies on humans or animals have been carried out, as well as in clinical studies, information about obtaining the consent of the Ethics Committee should be included.
- The Editorial Board follow the principles contained in the Helsinki Declaration as well as in the Interdisciplinary Principles and Guidelines for the Use of Animals in Research, Testing and Education, published by the New York Academy of Sciences Ad Hoc Committee on Animal Research. All papers relating to animals or humans must comply with ethical principles set out by the Ethics Committee.
- The abstract should contain 150-250 words. Abstracts of original, both clinical and experimental, papers should have the following structure: Aims, Material and methods, Results, Conclusions. Do not use abbreviations in the title or the abstract. The abstract is pasted or rewritten by the authors into the appropriate field in the application form in the editorial panel.
- Keywords (3-5) should be given according to MeSH (Medical Subject Headings Index Medicus catalogs – <http://www.nlm.nih.gov/mesh/MBrowser.html>). Keywords cannot be a repetition of the title of the manuscript.
- Illustrative material may be black and white or color photographs, clearly contrasting or drawings carefully made on a white background. With the exception of selected issues, the Journal is printed in shades of gray (black and white illustrations).
- The content of the figures, if present (e.g. on the charts), should also be in English
- Links to all tables and figures (round brackets) as well as references (square brackets) the author must place in the text of the article.
- Only references to which the author refers in the text should be included in the list of references ordered by citation. There should be no more than 30 items in original papers and no more than 40 items in review papers. Each item should contain: last names of all authors, first letters of first names, the title of the manuscript, the abbreviation of the journal title (according to Index Medicus), year, number, start and end page. For book items, please provide: author's (authors') last name, first letter of the first name, chapter title, book title, publisher, place and year of publication. It is allowed to cite websites with the URL and date of use of the article, and if possible the last names of the authors. Each literature item should have a reference in the text of the manuscript placed in square brackets, e.g. [1], [3-6]. Items should be organized as presented in Annex 1 to these Regulations.
- When submitting the article to the editor, the authors enclose a statement that the work was not published or submitted for publication in another journal and that they take full responsibility for its content, and the information that may indicate a conflict of interest, such as:
 - financial dependencies (employment, paid expertise, consulting, ownership of shares, fees),
 - personal dependencies,
 - academic and other competition that may affect the substantive side of the work,
 - sponsorship of all or part of the research at the stage of design, collection, analysis and interpretation of data, or report writing.
- The authors in the editorial panel define their contribution to the formation of scientific work according to the following key:
 - Work concept and design
 - Data collection and analysis
 - Responsibility for statistical analysis
 - Writing the article
 - Critical review
 - Final approval of the article.
- In the editorial panel along with the affiliation, the author also gives her or his ORCID number.
- The Journal is reviewed in double, blind review mode. The submitted papers are evaluated by two independent reviewers and then qualified for publishing by the Editor-in-Chief. Reviews are anonymous. The authors receive critical reviews with a request to correct the manuscript or with a decision not to qualify it for publishing. The procedure for reviewing articles is in line with the recommendations of the Ministry of Science and Higher Education contained in the paper "Good practices in review procedures in science" (Warsaw 2011). Detailed rules for dealing with improper publishing practices are in line with COPE guidelines. The publishing review rules are in the Review Rules section.
- Each manuscript is subject to verification in the anti-plagiarism system.
- Manuscripts are sent for the author's approval. The author's corrections should be sent within the time limit indicated in the system. No response within the given deadline is tantamount to the author's acceptance of the submitted material. In special cases, it is possible to set dates individually.
- Acceptance of the manuscript for publishing means the transfer of copyright to the Aluna Publishing House (Aluna Anna Łuczynska, NIP 5251624918).
- Articles published on-line and available in open access are published under Creative Commons Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0) allowing to download articles and share them with others as long as they credit the authors and the publisher, but without permission to change them in any way or use them commercially.
- The authors receive a free PDF of the issue in which their manuscript is enclosed, and on request – a printed copy. The printed copy is sent to the address indicated by the authors as the correspondence address.
- Manuscripts not concordant with the above instructions will be returned to be corrected.
- The editors do not return papers which have not been commissioned.
- The editors take no responsibility for the contents of the advertisements.

CONTENTS

ORIGINAL ARTICLES

- Aidyn G. Salmanov, Oleg M. Ishchak, Serhii A. Dobarin, Olena M. Susidko, Olena V. Mosendz, Svitlana M. Korniyenko, Oleksandr A. Voloshyn
PERINATAL INFECTIONS IN UKRAINE: RESULTS OF A MULTICENTER STUDY 2025
- Kateryna Hlushko, Oksana Boyarchuk, Maria Kinash, Emillia Burbela, Yana Rohalska, Lesia Dobrovol'ska
AWARENESS OF FOLIC ACID USE AND ITS EFFECTS AMONG MEDICAL STUDENTS IN UKRAINE 2033
- Iana O. Andreieva, Olha I. Riznyk, Sergii P. Myrnyi, Nikolai N. Surmylo
STATE OF CUTANEOUS MICROCIRCULATION IN PATIENTS WITH OBESITY 2039
- Bohdan Tataryn, Anna Kryzhanivska, Iryna Dyakiv, Alina Andriiv
SURVIVAL OF PATIENTS WITH RECTAL CANCER 2044
- Alina O. Ponyrko, Valentyna I. Bumeister, Serhii M. Dmytruk, Olha S. Yarmolenko, Tetiana P. Teslyk, Tetiana V. Riabenko, Iryna Shkolna
STRUCTURAL CHANGES OF LONG TUBULAR BONES OF MATURE RATS UNDER THE HYPERGLYCEMIA 2052
- Tetiana V. Harapko, Lesia R. Mateshuk-Vatseba
MORPHOMETRIC AND ELECTRON MICROSCOPIC CHANGES OF STRUCTURAL COMPONENTS OF LYMPH NODES DURING CORRECTION OF THE ACTION OF SODIUM GLUTAMATE WITH ORLISTAT 2060
- Inna I. Torianyk
ULTRAMICROSCOPIC CHARACTERISTICS OF ERYTHROCYTES IN INDIVIDUALS WITH BORRELIOSIS, RHEUMATOID ARTHRITIS AND TOXOPLASMOSIS AS A MARKER, COMORBID OR CONCOMITANT PATHOLOGY FOR BABESIOSIS 2066
- Kseniia Suprun, Yanina Kutasevych, Iryna Oliinyk, Oleksii Bilozorov, Iryna Mashtakova, Olga Stuliy
PECULIARITIES OF TREATMENT OF PATIENTS WITH ONYCHOMYCOSIS WITH CONTRAINDICATIONS TO SYSTEMIC ANTIMYCOTICS 2070
- Iryna M. Nikitina, Nataliia Ye. Horban, Kseniia V. Mykytyn, Kateryna O. Kondratiuk, Svitlana A. Smiian, Nataliia V. Kalashnyk, Nadiya V. Gerevich
THE STATE OF REPRODUCTIVE HEALTH OF WOMEN WITH HYPERPROLIFERATIVE PATHOLOGY OF THE ENDOMETRIUM 2076
- Olga L. Skrypa
EVALUATION OF THE EFFECTIVENESS OF TREATMENT OF PATIENTS WITH TEMPOROMANDIBULAR JOINT DYSFUNCTION AFTER MANDIBULAR FRACTURES ACCORDING TO X-RAY EXAMINATION 2082
- Igor P. Vakaliuk, Nataliia V. Savchuk, Roksolana V. Nesterak, Haliia B. Kulynych, Ruslana S. Hryhoryshyn
INFLUENCE OF COMPLEX TREATMENT WITH MAGNESIUM AND POTASSIUM SALTS OF GLUCONIC ACID, EPLERENONE AND RIVAROXABAN ON DYNAMICS OF INDICATORS OF ISCHEMIA AND MYOCARDIAL REMODELING IN PATIENTS WITH CHRONIC HEART FAILURE AFTER MYOCARDIAL INFARCTION 2087
- Oksana Ishchenko, Iryna Koshova, Inna Borysova, Dmytro Stepanskyi
MICROBIOLOGICAL FEATURES OF *STAPHYLOCOCCUS AUREUS* ISOLATED FROM RESPIRATORY TRACT OF CHILDREN WITH CYSTIC FIBROSIS 2094
- Petro A. Hasiuk, Mykhailo Ramus, Anna Vorobets, Iryna Tkachenko, Tetiana Dzetsiukh, Dmytro Kalashnikov, Svitlana Rosolovska
COMPARATIVE EVALUATION OF HISTOTOXICITY INDICATORS OF METAL ALLOYS FOR THE MANUFACTURE OF METAL CERAMIC DENTAL CONSTRUCTIONS 2100
- Antonina Sydorenko, Ruslan Lutsenko, Iryna Kniazkova, Nataliia Liakhovska, Olga Lutsenko
PECULIARITIES OF MOTOR ACTIVITY OF RATS IN SIMULATION PHARMACOLOGICAL DEPRESSOGENIC STATES 2105
- Evheniia A. Shtaniuk, Oleksandra O. Vovk, Larisa V. Krasnikova, Yuliia I. Polyvianna, Tetiana I. Kovalenko
STUDY OF ANTIBACTERIAL ACTIVITY OF THE PREPARATIONS WITH DIOXIDINE AND LEVOFLOXACIN ON MAIN PURULENT-INFLAMMATORY PROCESSES PATHOGENS 2109
- Natalia N. Brailko, Iryna M. Tkachenko, Victor V. Kovalenko, Anna V. Lemeshko, Alexey G. Fenko, Ruslan V. Kozak, Dmitry V. Kalashnikov
INVESTIGATION OF STRESS-STRAIN STATE OF "RESTORATION & TOOTH" SYSTEM IN WEDGE-SHAPED DEFECTS BY COMPUTED MODELING METHOD 2112
- Yuliya V. Sarkisova, Viktor T. Bachynskiy, Oleksandr I. Garazdiuk, Ivan V. Garazdiuk, Iana M. Teleki
POSTMORTEM INTERVAL ESTIMATION BY THE METHOD OF WAVELET ANALYSIS OF STOKES-POLARIMETRIC MAPPING DATA OF HUMAN VITREOUS BODY LAYERS 2118
- Volodymir K. Likhachov, Viktoriya L. Vashchenko, Olena O. Taranovska
IMPACT OF PREVENTIVE THERAPY ON REGULATING MECHANISMS OF DECIDUAL MACROPHAGE POLARIZATION IN PREGNANT WOMEN WITH HIGH RISK OF PREECLAMPSIA 2123
- Zbigniew Żęgota, Joanna Goździk, Joanna Głogowska-Szeląg
IMPROVED PHYSICAL FUNCTION WITH COMPLEMENTARY USE OF A DIETARY SUPPLEMENT FOR MILD KNEE OSTEOARTHRITIS: A SUBGROUP ANALYSIS 2128

Valentyna Sklyarova, Kateryna Shatylovich, Pavlo Sklyarov, Anzhelika Filipyuk SHOULD ASCARIASIS BE CONSIDERED AS A REPRODUCTOLOGY PROBLEM?	2138
Oleksandr Belov, Yevhen Loiko, Iryna Mazaikina, Iryna Pototska FEATURES OF PSYCHOPATHOLOGICAL REACTION IN PARENTS WHOSE CHILDREN SUFFER FROM CHRONIC MUSCULOSKELETAL DISORDERS	2147
Marta M. Zelinka-Khobzey, Kostiantyn V. Tarasenko, Tetiana V. Mamontova, Oksana A. Shlykova CHARACTERISTICS OF CD68+ AND CD163+ EXPRESSION IN PLACENTA OF WOMEN WITH PREECLAMPSIA AND OBESITY	2152
Yaroslav P. Feleshtynskiy, Sergiy O. Oparin, Bogdan V. Sorokin, Margaryta G. Boiarskaia, Dmytro V. Lutsenko ENDOSCOPIC HEMOSTASIS IN ULCERATIVE GASTRODUODENAL BLEEDING USING HIGH-FREQUENCY BIOLOGICAL WELDING ELECTROLIGATION	2159
Oksana V. Tymoshchuk, Zhanna Sotnikova-Meleshkina, Olha P. Melnyk-Sheremeta, Oksana V. Zahorodnieva, Krystyna A. Kryvonos, Kseniia M. Ostrovska KEY ASPECTS OF ORGANIZATION OF THE PROCESS AND INDIVIDUAL CHARACTERISTICS OF NUTRITION OF PUPILS AND STUDENTS OF DIFFERENT TYPES OF MODERN EDUCATIONAL INSTITUTIONS	2163
Nataliya O. Gutorova, Vitalii M. Pashkov, Oleksii S. Soloviov ILLEGAL INTERNET PHARMACIES AS A THREAT TO PUBLIC HEALTH IN EUROPE	2169
Natalya A. Maruta, Marianna V. Markova, Hanna M. Kozhyna, Tetiana A. Aliieva, Lyudmyla M. Yuryeva, Tsira B. Abdryakhimova, Nataliia G. Pshuk, Andrii M. Skrypnykov PSYCHOLOGICAL FACTORS AND CONSEQUENCES OF PSYCHOSOCIAL STRESS DURING THE PANDEMIC	2175
Igor S. Brodetskyi, Vladislav A. Malanchuk, Olena O. Dyadyk, Yaroslava A. Kulbashna, Olena O. Astapenko, Nataliia V. Kapustnyk, Mykhailo S. Myroshnychenko MORPHOLOGICAL JUSTIFICATION OF THE SALIVARY GLAND TISSUE RESECTION BOUNDARIES DURING SURGICAL TREATMENT OF PATIENTS WITH PLEOMORPHIC ADENOMAS	2182
Yurii Yu. Yarov, Iryna I. Tkachenko DYNAMICS OF ANTI-ANTIOXIDANT SYSTEM INDICATORS IN THE POSTOPERATIVE PERIOD IN PATIENTS WITH PERIODONTITIS ACCOMPANIED BY DIFFERENT REACTIVITY OF THE ORGANISM	2187
Vladislav V. Liubchak, Liliia M. Khomenko, Michael P. Kovalishyn, Viktoriia V. Ilyina, Tetiana V. Babar, Volodymyr V. Sikora, Olha V. Simonova NEW MARKETING TOOLS IN THE BLOOD SERVICE TO OPTIMIZE QUALITY MANAGEMENT	2192
Nazar R. Hrytsevych, Volodymyr V. Vereschaka FEATURES OF THE ULTRASTRUCTURE OF THE SKIN OF WHITE RATS 60 AND 90 DAYS AFTER MODELING OF PORTAL HYPERTENSION	2197
Roman Ye. Bulyk, Vladyslav R. Yosypenko, Kateryna V. Vlasova ONTOGENETIC ASPECTS OF MELATONIN RECEPTORS 1A DENSITY IN THE LATERAL PREOPTIC NUCLEUS OF THE HYPOTHALAMUS UNDER VARIOUS LIGHT CONDITIONS	2202
REVIEW ARTICLES	
Vitalii Pashkov, Maryna V. Trotska, Liudmyla M. Nikolenko AMBIENT AIR PROTECTION IN THE SYSTEM OF PUBLIC HEALTH AND BIOLOGICAL SECURITY LEGAL PROVISION	2207
Liudmyla A. Vygivska, Lesia A. Rudenko, Evgeniia B. Radzishavska, Iryna M. Merenkova, Violeta B. Kalnytska PECULIARITIES OF THE PREVALENCE OF INTRAUTERINE INFECTIONS AT THE PRESENT STAGE	2213
Anna Krakowiak, Marek Rajzer, Mateusz Gaczoł, Urszula Gancarczyk, Paweł Prochownik, Natalia Podolec, Zuzanna Sachajko, Filip Baranowski, Anna Pyczek, Monika Komar OBESITY AND ATRIAL FIBRILLATION – BARIATRIC SURGERY AS A METHOD OF AF RISK DECREASE	2218
Halyna V. Bilavych, Jan V. Bilawicz, Larysa V. Slyvka, Mariia V. Klepar, Mykola P. Pantiuk, Nadiya O. Fedchyshyn, Borys P. Savchuk SANITARY AND HYGIENIC EDUCATION OF CHILDREN AND ADULTS UNDER THE CONDITIONS OF THE COVID-19 PANDEMIC IN MEASUREMENT OF THE ACHIEVEMENTS OF PUBLIC MEDICINE OF GALICIA (END OF THE XIX – 30S OF THE XX CENTURY)	2222
Waldemar Gniadek MIECZYŚLAW KONOPACKI – THE OUTSTANDING POLISH HISTOLOGIST, ANATOMIST, AND FREEMASON ACTIVIST	2228
CASE STUDIES	
Vladimir Belokonev, Andrei I. Gritsaenko, Alexey Nikolaev, Tatiana Larina A CLINICAL CASE OF COEXISTENT PARASITIC LIVER AND GASTRIC LESIONS CAUSED BY CLONORCHIASIS	2232
Aleksandra Oraczewska, Marek Ochman, Mariola Ograbek-Król, Wiktoria Kowalska, Klaudia Glinka, Fryderyk Zawadzki, Adam Barczyk, Szymon Skoczyński PULMONARY ALVEOLAR MICROLITHIASIS. DISCREPANCIES BETWEEN RADIOLOGICAL FINDINGS AND CLINICAL PATTERN- CASE STUDY	2235

ORIGINAL ARTICLE

PSYCHOLOGICAL FACTORS AND CONSEQUENCES OF PSYCHOSOCIAL STRESS DURING THE PANDEMIC

DOI: 10.36740/WLek202109126

Natalya A. Maruta¹, Marianna V. Markova², Hanna M. Kozhyna³, Tetiana A. Aliieva², Lyudmyla M. Yuryeva⁴, Tsira B. Abdryakhimova⁵, Nataliia G. Pshuk⁶, Andrii M. Skrypnikov⁷

¹STATE INSTITUTION "INSTITUTE OF NEUROLOGY, PSYCHIATRY AND NARCOLOGY OF THE NAMS OF UKRAINE", KHARKIV, UKRAINE

²KHARKIV MEDICAL ACADEMY OF POSTGRADUATE EDUCATION, KHARKIV, UKRAINE

³KHARKIV NATIONAL MEDICAL UNIVERSITY, KHARKIV, UKRAINE

⁴STATE INSTITUTION "DNIPROPETROVSK MEDICAL ACADEMY OF THE MINISTRY OF HEALTH OF UKRAINE", DNIPRO, UKRAINE

⁵BOGOMOLETS NATIONAL MEDICAL UNIVERSITY, KYIV, UKRAINE

⁶NATIONAL PIROGOV MEMORIAL MEDICAL UNIVERSITY, VINNYTSIA, UKRAINE

⁷POLTAVA STATE MEDICAL UNIVERSITY, POLTAVA, UKRAINE

ABSTRACT

The aim: Study of psychological factors and consequences of psychosocial stress which is formed during the COVID-19 pandemic.

Materials and methods: In the research, we used methods: developed a common, assessment of psychosocial stress L. Reeder scale, assessment of distress R.Kessler, assessment of the presence of manifestations of anxiety-depressive response GAD-7, depression self-assessment scale PHQ-9, method of assessing neuropsychological adaptation I. Gurvich, assessment of psychosocial support D. Ziemet, the scale of assessment of family anxiety E. Eidemiller, W. Justickis, strategies for stress-coping behavior E. Heim, methods of assessing the quality of life A.A. Pushkarev, N.G. Arinchina (2000). The current factors of the impact of coronavirus on the mental state of the population have been investigated. The study involved 823 citizens of Ukraine, who filled out the developed Google form.

Results: Threats of coronavirus disease, violation of the usual life stereotype, restriction of leisure activity, harmful interest in news about the pandemic, usage of tobacco plays an important role in the mechanisms of distress formation. These conditions raise the risk of increased stress, anxiety and depression.

Conclusions: The investigations carried out suggested that the quarantine restrictions could be predisposing factors for mental health impairments. Under these conditions, risks of increased stress pressure, anxiety, and depression are rising.

Measures on psycho-prevention should be performed on the base of the regularities identified.

KEY WORDS: pandemic (COVID-19), psycho-emotional state, anxiety, stress, coping strategies, the quality of life.

Wiad Lek. 2021;74(9 p.1):2175-2181

INTRODUCTION

The 2019 coronavirus pandemic (COVID-19) has become a great challenge for people's mental health around the world.

High anxiety, depression, post-traumatic disorder and psychological distress rates are recorded among the nations during the pandemic [1-5].

Detection of the direct effect of COVID-19 on mental processes is a difficult problem, however, certain determinants identified in the conducted researches suggest the following [6]. Despite the undeniable need for isolation and physical distancing, which resist the high contagiousness and pathogenicity of the virus, those factors have a negative impact on mental health [7-10].

Research also notes the pathogenic role of psychosocial factors, including economic difficulties and financial loss (due to unemployment and declining incomes), school closures, lack of resources for health care, domestic violence and insufficient distribution of basic necessities.

Psychological consequences include fear of getting infected with COVID-19 and dying, loss of livelihood and loved ones, insecurity and anxiety about the future, social discrimination, separation from family and caretakers [11-14].

Common risk factors for mental disorders during the COVID-19 pandemic include women, the younger age group (≤ 40 years), the presence of chronic / mental distresses, unemployment, students and frequent conversations via social media.

There is a higher prevalence of adverse psychological effects in the population compared to the spread of the pandemic [15-17].

Symptoms of adverse psychological consequences were more common at the onset of the outbreak, when people were subjected to mandatory quarantine, sudden unemployment, and the uncertainty associated with the COVID-19 outbreak [18].

The duration of mental symptoms should also be considered when assessing the psychological consequences of a coronavirus outbreak, as acute psychological responses to stressful or traumatic events have sometimes a protective and evolutionary nature [19-20].

Several prognostic factors have been identified in the course of research. Women were more vulnerable to developing symptoms of various forms of mental disorders during a pandemic, including depression, anxiety, post-traumatic stress disorder, and adaptation disorders [21-23]. High severity of psychological stress occurs in women because they make up a higher percentage of the workforce, which can be negatively affected by COVID-19, for example, in retail, services and health care.

People under the age of 40 also had more adverse psychological symptoms during a pandemic [24-25]. This conclusion may be partly related to their role in caring for families (ie, especially women) that provide financial and moral support to children or the elderly. Job losses and unpredictability, caused by the COVID-19 pandemic, among this age group can be particularly stressful. In addition, the majority of people under the age of 40 are students who may also experience stronger emotional upheavals due to school closures, cancellation of public events, lower learning efficiency in remote online courses and postponement of exams [24].

These data conform the results testifying that student status was associated with higher levels of depressive symptoms and post-traumatic stress symptoms during the COVID-19 outbreak [27].

People with chronic diseases (somatic / mental) develop more symptoms of anxiety and stress during COVID-19 pandemic [28-30].

Anxiety and worrying about coronavirus infection in patients with chronic diseases is partly due to their weakened immunity caused by pre-existing conditions, which makes them susceptible to infection and a higher risk of death. Significantly higher mortality rates have been reported in patients with diabetes, hypertension and coronary heart disease, but the exact causes remain unknown. In addition, another practical aspect of concern for patients with pre-existing conditions may be the delay and unavailability of medical services and treatment as a result of the COVID-19 pandemic. For example, because the rapidly growing number of patients with COVID-19 used hospital and medical resources, this could inadvertently affect the primary, secondary, and tertiary prevention of other diseases. Individuals with a history of mental disorders or current diagnoses of mental illness are also usually more sensitive to external stressors, such as the social isolation associated with a pandemic [31].

A few researches have found that frequent use of social networks / news related to COVID-19 caused symptoms of anxiety and stress [32].

Frequent use of social networks, the influence of potentially fakes, misinformation are the factors of increasing concern. Due to the unpredictable situation and a lot of unknown information about the new coronavirus,

misinformation and fake news spread with ease through social networking platforms, creating unnecessary fears and concerns. Sadness and anxiety can also arise from the constant observation of members of a community suffering from a pandemic through social networking platforms or news reports [33].

THE AIM

The aim of our study was to identify the factors and psychological consequences of psychosocial stress, which is formed during the COVID-19 pandemic.

MATERIALS AND METHODS

In the study we used the following set of methods. We developed a common poll aimed at studying social demographic data, life conditions and lifestyle during the quarantine, the presence of chronic diseases and psychodiagnostic techniques (assessment of psychosocial stress) (L. Reeder scale), assessment of distress (R. Kessler scale), assessment of the presence of manifestations of anxiety-depressive response (GAD-7), depression self-assessment scale (PHQ-9), method of assessing neuropsychological adaptation (I. Gurvich), assessment of psychosocial support (D. Ziemet), the scale of assessment of family anxiety (E. Eidemiller, W. Justickis), strategies for stress-coping behavior (E. Heim), methods of assessing the quality of life (AA Pushkarev, NG Arinchina, 2000), as well as mathematical methods of data processing.

The study involved 823 people who filled out a Google survey form.

RESULTS

The analysis of the socio-demographic condition of the respondents showed the following.

The vast majority of respondents lived in the city – 88.21%; 6.77% – in settlements; 5.35% – in urban settlements. Residents of 20 regions of Ukraine (Vinnytsia, Volyn, Dnipro, Donetsk, Zhytomyr, Zakarpattia, Zaporizhia, Ivano-Frankivsk, Kyiv, Krapivnitsky, Lviv, Mykolaiv, Odessa, Poltava, Rivne, Sumy, Ternopil, Kharkiv, Khmelnytsky, Kherson, Chernivtsi) took part in the survey. The most represented among the participants were respondents from Kharkiv (21.38%), Dnipro (17.01%), Lviv (16.77%), Zaporizhia (10.08%) and Poltava (8.74%) areas.

The majority of the respondents were women (612 people, 74,36 %), men formed 25,64% (211 people). Analysis of the age characteristics of the respondents showed that the majority of respondents were under the age of 30 – 40.34% and aged 30-40 – 21.51%. At the age of 45-50 years there were 15.79% of respondents; at the age of 51-60 years – 12.27%; at the age of more than 60 years – 10.00%.

The marital status analysis showed that 47.51% were married, 43.5% – single, 8.99% – divorced. 51.03% of respondents did not have children; 23.57% had one child; 21.15% – two children; 4.25% – 3 and more children.

Analyzing the level of education, we noticed that 65.74%




Fig. 1. Lifestyle in the group examined before and during quarantine.




Fig. 2. The presence of chronic diseases in the group of subjects

of those surveyed had higher education; 12.76% – incomplete higher; 12.64% – average; 8.50% – special secondary; 0.36% – technical.

Living conditions in 56.62% were good; in 39.85% – satisfactory; in 3.52% – unsatisfactory.

In general, among the surveyed young people (up to 40 years old) predominated; female, childless; those with higher education and good living conditions.

During the study, we paid special attention to the analysis of living activity during the pandemic. The results showed that 60.14% of respondents comply with quarantine requirements and 39.86% – no. 24.54% continued to go to work during quarantine; 75.46% of respondents stayed at home (35.35% continued to work remotely).

In 37.67% of those surveyed, the work was related to the possibility of infection with COVID-19, in 62.33% this possibility was not detected.

Respondents’ assessments of their condition during quarantine were mixed. 74.24% of respondents believed

that they would “survive”; 52.36% said they felt “not very well”; 30.86% reported that they felt “terrible, I can’t stand the conditions of quarantine”; 17.25% reported the presence of a “feeling of loneliness.”

At the same time, 20.53% of respondents rated their condition as “good, similar to being on vacation”, and 17.49% expressed a desire to “extend quarantine”.

Studying the lifestyle before and during the quarantine period, it was found that the presence of excessive work before quarantine was found in 42.3%, and in quarantine – in 26.8% of respondents ($p \leq 0.05$); lack of permanent work was registered in 22.7% and 38.2%, respectively ($p \leq 0.05$) (Fig. 1).

Busy lifestyle was detected in 53.6% before and in 32.4% during quarantine (at $\varphi_{emp} = 8.804$ $\varphi_{cr} = 2.31$, where $p \leq 0.01$); 51.2% – noted the presence of work stress before, and 41.7% – during quarantine (at $\varphi_{emp} = 3.875$ $\varphi_{cr} = 2,31$, where $p \leq 0,01$); 32.7% – found a state of vigor to, and 39.8% – during quarantine (at $\varphi_{emp} = 3.367$ $\varphi_{cr} = 2.31$, where $p \leq 0.01$).

Summarizing the data obtained on the impact of quarantine on lifestyle, it should be noted that the number of overworked people in a stressful mode and in a hurry during quarantine decreased. One third of respondents (30.86%) reported distress during quarantine. Under quarantine conditions, the number of people who do not have a permanent job has increased.

Analysis of the presence of chronic diseases showed that most often the respondents had diseases of the gastrointestinal tract (18.71%), cardiovascular system (16.28%), respiratory system (11.78%), thyroid gland (11.66%) and neurological diseases (11.17%) (Fig. 2).

When studying the causes of anxiety (possible factors of mental trauma), the most important participants in the survey were selected:

- risk to life and health of relatives and friends (82.26%);
- impossibility of a habitual way of life (travel, sports, visiting theaters, museums) (79.82%);
- uncertainty in the actions of the authorities (79.10%);
- restrictions on leisure outside the home (74.11%);
- COVID-19 disease of family members, relatives, acquaintances (75.45%);
- lack of special treatment for COVID-19 (71.08%);
- the possibility of contracting a life-threatening disease (63.54%);
- danger to one's own life (63.18%).

Analyzing the presented data, it should be noted, that the vast majority of respondents were concerned about the risk to the life and health of loved ones, disruption of the usual way of life, restriction of leisure outside the home and uncertainty in the actions of the authorities.

Assessing the impact of news related to the COVID-19 pandemic, it was found that 44.85% of respondents considered it necessary to reduce the time spent searching for and reading such news; 17.25% felt guilty for searching and reading them; 25.15%, on the other hand, wanted to search and read pandemic news.

Assessing the level of interest in information about COVID-19, 51.40% of respondents considered this level to be normative; 9.60% noted interest in this information; 16.4% – risky interest, 22.60% – pathological (harmful) interest.

Analysis of alcohol consumption during quarantine showed that 13.24% of respondents had a feeling about the need to reduce alcohol consumption; 10.05% felt guilty about drinking alcohol; 8.75% were irritated when someone told them about the need to reduce alcohol consumption, 6.68% expressed a desire to drink alcohol, when they woke up.

Most of the respondents noted the absence of problems with alcohol (89.06%); 4.50% reported drinking alcohol; 2.67% – systematic alcohol consumption; 3.77% – alcohol dependence.

In the study of smoking, it was noted that tobacco use was registered in 22.60% of respondents. At the same time 9.11% burned the first cigarette within the first 5 minutes after waking up, 15.43% could give up the first cigarette; 11.78% burned 10 or fewer cigarettes a day (5.58% burned 11-20 cigarettes; 2.79% burned 21-30 cigarettes). 13.12% of

respondents smoked more often in the early hours of the morning; 7.04% reported continuing to smoke when ill. Assessment of the effects of smoking showed that tobacco dependence was not found in 40.80%; weak or moderate tobacco dependence was registered in 44.62%; strongly expressed dependence – in 14.52%.

That is, when assessing the informational effects, the effects of alcohol and tobacco during quarantine, it should be noted the presence of harmful interest in news about the pandemic (22.6%); low rates of systematic use (2.67%) and alcohol dependence (3.77%); high rates of weak and moderate dependence on tobacco (44.62%), as well as strong dependence (14.52%).

Alcohol before quarantine was consumed by 13.6% of respondents, during quarantine – 12.4%; tobacco smoking was reported by 25.8% before and 38.7% during quarantine.

Given the huge role of social support in the genesis of social stress, we conducted a study of family anxiety. According to the results of such a study, it was found that 31.71% of respondents showed a sense of guilt, 59.41% had a high level of anxiety, 62.21% had a high level of stress. A high overall rate of family anxiety was found in 68.53%.

It should be noted that according to the multidimensional scale of perception of social support (MSPSS), the respondents showed a high level of this indicator in terms of support for family (82.74%), friends (82.50%), significant people (70.23%) and overall indicator of the level of support (89.42%).

Summarizing the data on the emotional load of the respondents, it should be noted that the vast majority of respondents felt anxiety of a minimal and moderate level; medium and high levels of anxiety were registered in 18.46% and 8.26%, respectively. The indicators of the level of depression were similar: in the vast majority of respondents the level of depression was minimal, mild and moderate, severe and very severe depression was found in 11.91% and 5.22%, respectively. Despite the high level of social support, 68.53% showed a high overall rate of family anxiety.

A study of the level of anxiety using GAD7 showed that the minimum anxiety was determined by 43.25%, moderate – 30.01%, medium – 18.46%, high – 8.26%.

Analyzing the level of depression, we noted that 36.21% had minimal depression, 30.01% – mild, 16.64% – moderate, 11.91% – severe, 5.22% – very severe.

Assessment of stress levels in a group of respondents according to the method of Kessler R.C. showed that most of the subjects had a low level of stress (80.19%), 18.46% – an average level, 1.34% – a high level of stress.

Assessing the level of stress, anxiety and depression in the respondents, it should be noted that the average level of stress was 18.47%, and high – in 1.34%.

18.47% had an average level of anxiety, and 8.26% – high. Severe depression was determined by 11.91%, very severe – 5.22%.

Respondents' quality of life was assessed in terms of physical health, emotional state, social and cognitive functions.

Table I. Quality of life assessment

	823 people		M ± m average value
	absolute number	%	
Physical mobility			
<i>low level (10-30 point)</i>	67	8,14	69,5±2,51
<i>average level (35-65 point)</i>	288	34,99	
<i>high level (65-100 point)</i>	468	56,87	
Emotional condition			
<i>low level (10-30 point)</i>	128	15,55	72,39±3,27
<i>average level (35-65 point)</i>	336	40,83	
<i>high level (65-100 point)</i>	359	43,62	
Social functions			
<i>low level (10-30 point)</i>	84	10,21	69,64±2,68
<i>average level (35-65 point)</i>	231	28,07	
<i>high level (65-100 point)</i>	508	61,72	
Cognitive function			
<i>low level (10-30 point)</i>	62	7,53	96,05±3,24
<i>average level (35-65 point)</i>	204	24,79	
<i>high level (65-100 point)</i>	557	67,68	

The majority of respondents assessed their physical health at a high level (56,87%), 34, 99% – at the average level, only 8.14% is at the low level. At the same time, a quantitative analysis of the results on separate scales showed an increase in indicators that points at a deterioration in health during daily exercise (5.64 points) and a discrepancy between the state of Health and the desire to acquire a profitable job (5.77 points) (Table I).

The analysis of the emotional component of the respondents' quality of life showed a predominance of high (43.62%) and average (40.83%) levels of the indicator. The presence of a low level was noted by 15.55% of respondents. The analysis of this indicator on the scales reflected high values on the scale of "I feel that no one cares about me" (5.65 points) and on the scale of "I feel oppressed and depressed" (5.32 points).

The study of social functions within the quality of life showed that 61.72% identified a high level of social activity, 28.07% – an average level, and 10.21% – a low level. Individual scales analysis showed high scores on the scales "my health condition interferes with normal vacation" (6.06 points).

During the studying of cognitive functions, it was found that 67.68% of the respondents noted the presence of a high level of cognitive functions, 24.79% – an average level; 7.53% – a low level. The highest scale in this section was "my health condition leads to financial problems" (6.04 points).

Summarizing the data obtained in this study, the following should be noted.

Socio-demographic characteristics of respondents (823 people) included the predominance of the female (74.36%), age groups up to 40 years (61.85%), city resi-

dents (88.21%), single and divorced (52.49%), people with higher education (65.74%), good and satisfactory living conditions (56.62% and 39.85%, respectively).

When studying the impact of quarantine restrictions on the lifestyle of respondents, it was found that the number of people working excessively in stressful and hasty mode during quarantine had decreased. At the same time, the number of people who do not have a permanent job has increased under quarantine conditions, 30.86% of those surveyed reported during quarantine.

Analysis of distress factors (causes of anxiety) showed that the overwhelming majority of respondents were worried under the influence of such factors as risk to the life and health of loved ones (82.26%), violation of the usual lifestyle (79.82%), uncertainty in the actions of the authorities (79.10%), restriction of leisure time outside the home (74.11%).

Assessing lifestyle changes during quarantine, it was detected that 22.6% of respondents noted a harmful interest in news about the pandemic; 44.62% – weak and moderate dependence on tobacco, low rates of systematic use and alcohol dependence (2.67% and 3.77%, respectively).

A study of stress, anxiety, and depression levels found that 18.47% had moderate stress levels and 1.3% had high levels. Moderate and high rates of anxiety and depression were recorded in 1/5 – 1/4 of the respondents. 18.46% had an average level of anxiety, and 8.26% had a high level of anxiety. The severe level of depression was determined by 11.91%, very high level by 5.22%.

The analysis of examination of the quality of life has demonstrated, that 8.14% had a low level of its physical component, 15.55% had a low level of its emotional component, and 10.21% had a low level of a social activity.

DISCUSSION

Summarizing the data obtained in this study, it should be noted that quarantine restrictions in some way affect the lifestyle of respondents.

It has been noted that during quarantine the number of people overworking in stress and haste mode is decreasing. At the same time, the number of people who have a permanent job is decreasing and the number of people who have distress during quarantine is increasing. The next pattern is the influence of distress factors, which manifested themselves in the form of concern for the lives and health of loved ones, concern for the violation of normal lifestyles, insecurity in actions of the authorities and restrictions on leisure outside. Also, we have found changes in lifestyle during quarantine, which manifested in the presence of harmful interest in the news and the presence of weak or moderate dependence, mainly on tobacco. Alcohol dependence in our study was less pronounced.

According to the results of the study of the level of stress, anxiety and depression, it has been found that the respondents have a medium and high level of anxiety and depression.

We obtained interesting data when assessing the quality of life of respondents. Despite the fact that the overall indicators of quality of life and performance in the physical, emotional, social and cognitive spheres were high, quantitative analysis of individual scales revealed the following negative trends. These are the deterioration of health during daily activities, the inconsistency of the state of health with the desire to get a well-paid job, feelings of loneliness and depression; impossibility to go on a good vacation and prevent financial problems due to the state of health. The data obtained in the study provide some prospects for psychoprophylaxis and psychoeducation among the population to strengthen the psychological state and mental health of the population.

CONCLUSIONS

Summarizing the data obtained in 823 examined patients during quarantine during the Covid-19 pandemic, we can describe the condition as follows:

1. The investigations carried out suggested that the quarantine restrictions could be predisposing factors for mental health impairments.
2. The threat of coronavirus disease, a disruption of a habitual life stereotype, leisure restrictions, a harmful interest in news about the pandemic, and tobacco abuse play an important role in mechanisms of distress formation.
3. Under these conditions, risks of increased stress pressure, anxiety, and depression are rising.
4. The analysis of examination of the quality of life has demonstrated, that 8.14 % had a low level of its physical component, 15.55 % had a low level of its emotional component, and 10.21 % had a low level of a social activity.
5. Measures on psycho-prevention should be performed on the base of the regularities identified.

REFERENCES

1. Wang C., Pan R., Wan X., Tan Y., Xu L., Ho C.S., Ho R.C. Immediate psychological responses and associated factors during the initial stage of the 2019 coronavirus disease (COVID-19) epidemic among the general population in China. *Int. J. Environ. Res. Public Health*. 2020;17(5):1729. doi: 10.3390/ijerph17051729.
2. Wasserman D, van der Gaag R, Wise J. The term “physical distancing” is recommended rather than “social distancing” during the COVID-19 pandemic for reducing feelings of rejection among people with mental health problems. *Eur Psychiatry*. 2020;63:e52.
3. World Health Organization (WHO) [Internet], Coronavirus disease (COVID-2019) situation reports [cited 2020 Aug 15]. Available from: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports>.
4. World Health Organization (WHO) [Internet], Mental health and COVID-19 [cited 2020 Aug 15]. Available from: <http://www.euro.who.int/en/health-topics/noncommunicable-diseases/mental-health/data-and-resources/mental-health-and-covid-19>.
5. World Health Organization (WHO) [Internet]. WHO briefing note – Mental health and psychosocial considerations during COVID-19 outbreak [cited 2020 Aug 15], Available from: <https://www.who.int/docs/default-source/coronaviruse/mental-health-considerations.pdf?sfvrsn=6d3578aM0>.
6. Karpenko Olga A., Syunyakov Timur S., Kulygina Maya A., et al. Impact of COVID-19 pandemic on anxiety, depression and distress – online survey results amid the pandemic in Russia. *Consortium Psychiatricum*. 2020; 1: 8-20.
7. Hao F, Tan W, Jiang L, et al. Do psychiatric patients experience more psychiatric symptoms during COVID-19 pandemic and lockdown? A case-control study with service and research implications for immunopsychiatry. *Brain Behav Immun*. 2020;87:100–6.
8. Pfefferbaum B, North CS. Mental health and the Covid-19 pandemic. *N Engl J Medicine*. 2020;383:510–2.
9. Vindegaard N, Eriksen Benros M. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brain Behav Immun*. 2020;S0889-1591:30954–5.
10. Zhang SX, Wang Y, Rauch A, Wei F. Unprecedented disruption of lives and work: health, distress and life satisfaction of working adults in China one month into the COVID-19 outbreak. *Psychiatry Res*. 2020;288:112958.
11. Adhanom Ghebreyesus T. Addressing mental health needs: an integral part of COVID-19 response. *World Psychiatry*. 2020;19:129–30.
12. Gruber C. Impaired interferon signature in severe COVID-19. *Nat Rev Immunol*. 2020;20:353.
13. Unützer J, Kimmel RJ, Snowden M. Psychiatry in the age of COVID-19. *World Psychiatry*. 2020;19:130–1
14. Wright L, Steptoe A, Fancourt D. Are we all in this together? Longitudinal assessment of cumulative adversities by socioeconomic position in the first 3 weeks of lockdown in the UK. *J Epidemiol Community Health*. 2020;74:683–8.
15. Huang Y. Prevalence of mental disorders in China: a cross-sectional epidemiological study. *Lancet Psychiat*. 2019;6(3):211–224. doi: 10.1016/S2215-0366(18)30511-X.
16. Lim G.Y., Tam W.W., Lu Y., et al. Prevalence of depression in the community from 30 countries between 1994 and 2014. *Sci. Rep*. 2018;8(1):2861. doi: 10.1038/s41598-018-21243-x.
17. Moghanibashi-Mansourieh A. Assessing the anxiety level of Iranian general population during COVID-19. *Asian J. Psychiatr*. 2020;51 doi: 10.1016/j.ajp.2020.102076.

18. Ho C.S.H., Chee C.Y., Ho R.C.M. Mental health strategies to combat the psychological impact of coronavirus disease (COVID-19) beyond paranoia and panic. *Ann. Acad. Med. Singapore*. 2020;49(3):155–160.
19. Gilbert P. Evolution and depression: issues and implications. *Psycho. Med.* 2006;36(3):287–297. doi: 10.1017/S0033291705006112.
20. Yaribeygi H., Panahi Y., Sahraei H., et al. The impact of stress on body function: a review. *EXCLI J.* 2017;16:1057–1072. doi: 10.17179/excli2017-480.
21. Ahmed M.Z., Ahmed O., Zhou A., et al. Epidemic of COVID-19 in China and associated psychological problems. *Asian J. Psychiatr.* 2020;51 doi: 10.1016/j.ajp.2020.102092.
22. Gao J., Zheng P., Jia Y., et al. Mental health problems and social media exposure during COVID-19 outbreak. *PLoS ONE*. 2020;15(4) doi: 10.1371/journal.pone.0231924.
23. Lei L., Huang X., Zhang S., et al. Comparison of prevalence and associated factors of anxiety and depression among people affected by versus people unaffected by quarantine during the covid-19 epidemic in southwestern China. *Med. Sci. Monit.* 2020;26 doi: 10.12659/MSM.924609.
24. Cao W., Fang Z., Hou G., et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res.* 2020;287 doi: 10.1016/j.psychres.2020.112934.
25. Huang Y., Zhao N. Generalized anxiety disorder, depressive symptoms and sleep quality during COVID-19 outbreak in China: a web-based cross-sectional survey. *Psychiatry Res.* 2020;288 doi: 10.1016/j.psychres.2020.112954.
26. Olagoke A.A., Olagoke O.O., Hughes A.M. Exposure to coronavirus news on mainstream media: the role of risk perceptions and depression. *Br. J. Health Psychol.* 2020 doi: 10.1111/bjhp.12427.
27. Samadarshi S.C.A., Sharma S., Bhatta J. An online survey of factors associated with self-perceived stress during the initial stage of the COVID-19 outbreak in Nepal. *Ethiop. J. Health Dev.* 2020;34(2):1–6.
28. Mazza C., Ricci E., Biondi S., et al. A nationwide survey of psychological distress among Italian people during the COVID-19 pandemic: immediate psychological responses and associated factors. *Int. J. Environ. Res. Public Health.* 2020;17:3165. doi: 10.3390/ijerph17093165.
29. Ozamiz-Etxebarria N., Dosil-Santamaria M., Picaza-Gorrochategui M., et al. Stress, anxiety and depression levels in the initial stage of the COVID-19 outbreak in a population sample in the northern Spain. *Cad. Saude. Publica.* 2020;36(4) doi: 10.1590/0102-311X00054020.
30. Özdin S., Özdin S.B. Levels and predictors of anxiety, depression and health anxiety during COVID-19 pandemic in Turkish society: the importance of gender. *Int. J. Soc. Psychiatry.* 2020:1–8. doi: 10.1177/0020764020927051.
31. Li Z. Vicarious traumatization in the general public, members, and non-members of medical teams aiding in COVID-19 control. *Brain Behav. Immun.* 2020 doi: 10.1016/j.bbi.2020.03.007.
32. Pierce M., McManus S., Jessop C., et al. Says who? The significance of sampling in mental health surveys during COVID-19. *Lancet Psychiatry.* 2020;7(7):567–8. Epub 2020/06/06. doi: 10.1016/s2215-0366(20)30237-6.
33. Erku D.A., Belachew S.W., Abrha S., et al. When fear and misinformation go viral: pharmacists' role in deterring medication misinformation during the 'infodemic' surrounding COVID-19. *Res. Social. Adm. Pharm.* 2020. doi: 10.1016/j.sapharm.2020.04.032.

The research was carried out within the framework of the scientific research work of the Department of Psychotherapy of the Kharkiv Medical Academy of Postgraduate Education of the Ministry of Health of Ukraine on the topic "Mental health disorders under social stress: clinical phenomenology, predictors of formation, system of psychodiagnostics, psychotherapy and psychological rehabilitation" (state registration number 0120U101490).

ORCID and contributionship:

Natalya A. Maruta: 0000-0002-6619-9150 ^{A,E,F}
 Marianna V. Markova: 0000-0003-0726-4925 ^{B,D,F}
 Hanna M. Kozhyna: 0000-0002-2000-707X ^{C,F}
 Tetiana A. Aliieva: 0000-0003-3511-027X ^{B,F}
 Lyudmyla M. Yuryeva: 0000-0002-1713-1037 ^{D,F}
 Tsira B. Abdryakhimova: 0000-0002-9406-2696 ^{C,F}
 Nataliia G. Pshuk: 0000-0001-9857-2565 ^{B,F}
 Andrii M. Skrypnykov: 0000-0002-1562-9421 ^{E,F}

Conflict of interest:

The Author declare no conflict of interest

CORRESPONDING AUTHOR

Natalya A. Maruta

Department of Borderline Psychiatry of the
 SI "Institute of Neurology, Psychiatry and
 Narcology of the NAMS of Ukraine"
 Akademika Pavlova Street, 46, Kharkiv, 61068, Ukraine
 tel: +380577255813
 e-mail: mscience@ukr.net

Received: 29.05.2021

Accepted: 25.08.2021

A – Work concept and design, B – Data collection and analysis, C – Responsibility for statistical analysis,
 D – Writing the article, E – Critical review, F – Final approval of the article