

VOLUME LXXVI, ISSUE 3, MARCH 2023

ISSN 0043-5147  
E-ISSN 2719-342X

# Wiadomości Lekarskie Medical Advances



Official journal of Polish Medical Association has been published since 1928



INDEXED IN PUBMED/MEDLINE, SCOPUS, EMBASE, EBSCO, INDEX COPERNICUS,  
POLISH MINISTRY OF EDUCATION AND SCIENCE, POLISH MEDICAL BIBLIOGRAPHY

Olexii I. Dronov, Inna O. Kovalska, Yelyzaveta. S. Kozachuk, Liudmyla V. Levchenko, Dmytro A. Vlasenko, Andrii S. Shvets CHANGES ANALYSIS OF THE HEPATOCYTE APOPTOSIS MARKERS LEVELS IN MALIGNANT OBSTRUCTIVE JAUNDICE COMPLICATED BY CHOLANGITIS	560
Olena A. Dulo, Yurii M. Furman, Olha B. Maltseva, Svitlana M. Samoilenko PHYSICAL HEALTH OF FEMALES FROM THE LOWLAND DISTRICTS OF ZAKARPATTIA ACCORDING TO THE METABOLIC LEVEL OF AEROBIC AND ANAEROBIC ENERGY SUPPLY DEPENDING ON THE COMPONENT BODY COMPOSITION	568
Viktoriia Z. Ivaskevych, Anatoliy M. Potapchuk, Oleh Yu. Rivis, Mariya V. Rivis, Yuriy V. Rak, Roman Yu. Marukha THE DETERMINATION OF THE NEED TO PROVIDE ORTHODONTIC ASSISTANCE TO TEENAGERS IN CONDITIONS OF LIMITED RESOURCES	575
Vitalina V. Ivachevska, Mykhailo M. Ivachevskyi, Mykhailo M. Hechko, Ivan I. Myhovych, Olga S. Blaga EFFICACY OF COMPREHENSIVE TREATMENT OF NONALCOHOLIC FATTY LIVER DISEASE IN PATIENTS WITH PREDIABETES	581
Volodymyr M. Bilak, Lyudmila V. Ignatko, Natalya V. Sochka, Olena V. Debretseni, Gabriella B. Kossey, Volodymyr Y. Mashika, Taras I. Griadil THE INFLUENCE OF SPELEOTHERAPY ON BRONCHI PASSAGE IN CHILDREN WITH BRONCHIAL ASTHMA USING A PHARMACO-FUNCTIONAL TEST WITH SALBUTAMOL	586
Alexander N. Stoyanov, Serhii S. Mashchenko, Valeriy I. Kalashnikov, Rooslan S. Vastyanov, Alexander R. Pulyk, Tamara O. Andreeva, Olena O. Kolesnik VESTIBULAR DYSFUNCTIONS IN CHRONIC BRAIN ISCHEMIA IN THE POST COVID PERIOD	591
Svitlana Yu. Karatieieva, Oleksandr M. Slobodian, Natalya Ya. Muzyka, Kseniya V. Slobodian, Oksana V. Kolesnik THE DETERMINATION OF HIP CIRCUMFERENCE IN THE MIDDLE OF YOUNG BOYS AND YOUNG GIRLS OF HIGHER EDUCATION INSTITUTIONS OF BUKOVINA DEPENDING ON THE SPORT TYPE	597
Yaroslav M. Popovich, Myroslav V. Rosul, Paula R. Sich, Orest P. Laver THROMBOLYSIS IN PULMONARY EMBOLISM TREATMENT	604
Valerii V. Korsak, Yurii Y. Bobyk, Iryna I. Patskan OBSTETRIC AND PERINATAL ASPECTS OF METABOLIC DISORDERS IN PREGNANT WOMEN	610
Stepan S. Filip, Rudolf M. Slyvka, Andriy M. Bratasyuk, Yuriy P. Skripinets, Anatoliy I. Shitev EARLY DIAGNOSIS OF ASYMPTOMATIC CHRONIC ISCHEMIA OF THE LOWER EXTREMITIES	616
Yaroslav P. Feleshtynskyi, Oleh S. Marshtupa, Volodymyr F. Vatamaniuk DIFFERENTIATED CHOICE OF POSTERIOR METHODS OF DISCONNECTION OF ANATOMICAL COMPONENTS OF THE ABDOMINAL WALL IN COMBINATION WITH ALLOPLASTY IN POSTOPERATIVE VENTRAL HERNIAS OF GIANT SIZE	623
Roman M. Mitsoda, Kateryna-Mariya R. Mitsoda T-CRITERION AS A TOOL FOR DETERMINING THE RISK OF COMPLICATIONS OF THE GESTATIONAL PROCESS	629
Yelyzaveta S. Sirchak, Monika T. Maroshan, Yevheniia E. Dankanych, Olesia P. Balazh, Valentina Y. Koval BLOOD COAGULATION DISORDERS IN PATIENTS WITH LIVER CIRRHOSIS INFECTED COVID-19	634
Renata Yu. Pohorilyak, Andriya V. Zheliznyak, Olga V. Feger IMPACT OF DISTANCE EDUCATION ON STUDENTS' HEALTH	640

## ORIGINAL ARTICLE

## VESTIBULAR DYSFUNCTIONS IN CHRONIC BRAIN ISCHEMIA IN THE POST COVID PERIOD

DOI: 10.36740/WLek202303121

Alexander N. Stoyanov<sup>1</sup>, Serhii S. Mashchenko<sup>1</sup>, Valeriy I. Kalashnikov<sup>2</sup>, Rooslan S. Vastyanov<sup>2</sup>, Alexander R. Pulyk<sup>3</sup>, Tamara O. Andreeva<sup>4</sup>, Olena O. Kolesnik<sup>4</sup>

<sup>1</sup>ODESA NATIONAL MEDICAL UNIVERSITY, ODESA, UKRAINE

<sup>2</sup>KHARKIV NATIONAL MEDICAL UNIVERSITY, KHARKIV, UKRAINE

<sup>3</sup>STATE HIGHER EDUCATIONAL ESTABLISHMENT «UZHGOROD NATIONAL UNIVERSITY», UZHGOROD, UKRAINE

<sup>4</sup>BLACK SEA NATIONAL UNIVERSITY NAMED AFTER P. MOHYLA, MYKOLAIV, UKRAINE

### ABSTRACT

**The aim:** The aim of the study is the clinical-pathogenetic reasoning of vestibular dysfunctions (VD) development against the background of chronic brain ischemia in the presence of degenerative changes in the cervical spine (CS) in the post COVID period.

**Materials and methods:** 82 patients, in the conditions of the clinical base of the Odessa National Medical University in 2019-2021 were examined. Group I with VD against the background of chronic brain ischemia (CBI) at the compensated phase; Group II with VD against the background of CBI at the subcompensated phase (33 men; 49 women), aged from 18 to 55 years. The control group (CG) consisted of 20 patients of the corresponding gender and age. The condition of the state of the autonomic nervous system, vestibular functions, cervical spine, cerebral arteries and emotional condition were examined.

**Results:** Vestibulo-ataxic disorders were higher compared to CG and increased along with the degree of brain damage. An important aspect of the development of VD is autonomic dysfunction against the background of pathological autonomic characteristics with predominant parasympathetic orientation of autonomic tone, especially in the case of insufficiency of autonomic reactivity (AR) and pathological autonomic support of activity. Such changes significantly increased in the presence of subcompensation of CBI. The correlation between psychoemotional disorders and changes in autonomic characteristics with VD against the background of CBI with initial regularities depending on the degree of brain damage was defined. The progression of CBI is facilitated by coronavirus infection and manifested in autonomic and psychoemotional dysfunctions. A characteristic hemodynamic feature in groups with compensated and subcompensated CBI is the presence of reduced perfusion in basilar (BA) and vertebral (VA) arteries. Changes in cerebral vascular reactivity with a decrease in cerebrovascular reactivity indicators were characteristic of the subcompensated phase of CBI. Hyperactivity to rotational functional loads in both clinical groups has a high correlation with the presence of stair descent and, to a lesser extent, isolated instability in CS.

**Conclusions:** 1. The occurrence of VD is facilitated by the presence of autonomic dysfunction and degenerative-dystrophic changes in the CS, especially in case of subcompensation of CBI. 2. Psychoemotional changes were a characteristic feature of patients with VD against the background of CBI and had certain regularities depending on the phase of CBI. 3. Suffered coronavirus infection contributes to the progression of VD and further decompensation of CBI due to direct damage to the autonomic and vascular systems of the brain. 4. Changes in cerebral hemodynamics in the form of reduced perfusion in BA and VA, a decrease in cerebrovascular reactivity, and an increase in reactivity to rotational functional load were determined in patients with VD against the background of subcompensated CBI.

**KEY WORDS:** chronic brain ischemia, autonomic dysfunctions, degenerative-dystrophic damage, cervical spine, cerebrovascular reactivity

Wiad Lek. 2023;76(3):591-596

### INTRODUCTION

Chronic brain ischemia (CBI) in the structure of cerebrovascular pathology occurs much more often than acute conditions and leads to a long-term disability. It is the risk factor for the development of cerebral strokes. Prevention of the development of significant organic vascular changes in the brain is the most urgent problem of modern medicine and neurology [1-4]. The presence of vestibular dysfunctions (VD) in the clinical picture of vascular cerebral dyschemia correlates with the degree of

brain damage [5,6]. VD is the most common syndrome in ischemic brain damage. The clinical variety of dizziness occurs against the background of a decrease in the compensatory capabilities of the central mechanisms of balance, the sensory system, especially in case of CBI with a predominant lesion of the vertebral-basilar basin against the background of the cervical spine (CS) pathology [7]. The important indicator of VD occurrence is the change in vegetative and vascular reactivity, which complicates the examination and differential diagnosis

of dizziness. The progression of ischemic brain damage is advanced by the development of the coronavirus infection, which negatively impacts the process of neurological rehabilitation [8-14]. The penetration of SARS-CoV-2 virus into the central nervous system (CNS) causes its damage and can activate some degenerative nervous system diseases [15]. Hypercoagulation due to systemic inflammation, cytokine storm, endotheliitis and other factors lead to infectious vascular lesions of the CNS [16,17]. Inflammatory damage of the vestibular apparatus reduces blood flow, causes vasospasm, leads to thrombosis as well as other vascular problems that may occur during the coronavirus infection.

## THE AIM

The aim is the clinical-pathogenetic reasoning of VD development against the background of chronic brain ischemia in the presence of degenerative changes in the cervical spine (CS) in the post COVID period.

## MATERIALS AND METHODS

82 patients were examined, Group I with VD against the background of CBI at the compensated phase (n =24); Group II with VD against the background of CBI at the subcompensated (n=38) phase (33 men; 49 women), aged from 18 to 55 years, the average age was  $38.6 \pm 1.6$  years. All patients had a history of COVID 19. Control group (CG), n=20 – relatively healthy people who underwent professional screening. During the examination of the state of the autonomic nervous system (ANS) according to the questionnaire [8] all patients reported the dysfunction availability. In order to objectify statics and balance, an original electrical contact device was designed and used for ataxia assessment [18] with the calculation of the integrative index of ataxia (IIA). Emotional layers were examined using the hospital anxiety and depression scale [19], the cognitive sphere – according to the test of A.R. Luria [18]. All patients had an X-ray examination of the cervical spine with functional loads. X-rays of the cervical spine were performed using a digital radio-fluoroscopy system OPERA T90cex GMM (Italy). X-rays were executed in standard modes, in direct and lateral projections with functional load (flexion – extension).

Cerebral arteries were examined in triplex mode on an Ultima-PA ultrasound scanner (RADMYR, Ukraine). The parameters of time-averaged maximum blood flow velocity (TAMX) and resistance index (RI) in the anterior (ACA), middle (MCA), posterior (PCA) cerebral, vertebral (VA) and basilar (BA) arteries were studied. The state of cerebrovascular reactivity was assessed

using the following functional loads: hypercapnic reactivity (reactivity coefficient to hypercapnic (CrCO<sub>2</sub>)), and hyperventilation reactivity (CrO<sub>2</sub>) test, functional nitroglycerin (CrFNT) tests, functional probes with rotation (RIRP), index of the vasomotor reactivity (IVMR). Statistics 8.0 was used for statistical processing with the assessment of the Wilcoxon test, as well as the Mann-Whitney U test. Differences at  $p < 0.05$  were considered statistically significant.

## RESULTS

In the main groups, all patients complained of dizziness (predominantly non-systemic – 51.6%, systemic – 48.4%), caused by physical exertion (25.8%), head movements (38.7%), orthostatic changes (14.5 %), blood pressure increase (9.7%).

The main symptoms of the launch of accompanying pathological mechanisms of vestibulopathy have been revealed. Headache (75.8%) had vascular and/or autonomic components: vasomotor (24.2%), ischemic-hypoxic (48.4%), venous (27.4%) cephalgia. Localization: diffuse (40.3%), in the occipital (33.9%), parietal (14.5%), frontal (11.3%) regions. The algic syndrome was manifested by pain in the neck (69.3%), including the ones with irradiation to the shoulder and corresponding hand (37.1%); back pain (56.4%); cardialgias (37.1%) and other pain phenomena. Muscle tension of shoulder girdle and neck had reflex-tonic or generalized character in 48.4% of cases. In addition, the most significant symptoms were: increased blood pressure (59.8%), noise in the head (38.7%), decreased hearing (30.6%), orthostatic hypotension (19.3%), asthenia (41.9%), emotional lability (58.1%), cognitive disorders (70.9%), dyssomnias (40.3%). In the objective, vestibulo-ataxic disorders of mild or moderate severity prevailed [9, 13]. The average values of IIA were increased in patients with CBI compared to healthy subjects (Group I –  $2.7 \pm 0.09$  relative units, Group II –  $3.6 \pm 1.10$  relative units, CG –  $1.8 \pm 0.08$  relative units,  $p < 0.05$ ). High rates of IIA were accompanied by the presence of vagotonia ( $3.0 \pm 0.12$  relative units versus  $2.3 \pm 0.14$  relative units in patients with eytonia,  $p < 0.05$ ). Manifestations of cervical osteochondrosis were diagnosed in all patients. Instability of the cervical spine was noted in half of the examined patients, more often in segments C<sub>4</sub> – C<sub>5</sub> (77.4%,  $p < 0.05$ ), less often – C<sub>3</sub> – C<sub>4</sub> and C<sub>5</sub> – C<sub>6</sub>. Uncovertebral arthrosis was defined in 41.9% of studies. Pathological autonomic tone (AT) was manifested in 96.7% of patients, mostly shifted towards the vagal direction (53.2%); pathological autonomic reactivity (AR) – 83.9% of patients mainly had autonomic insufficiency (52.4%), which increased along with ischemic brain damage increase; changes in

**Table I.** Indexes TAMX (cm/s) in cerebral arteries in patients with CBI

	MCA	ACA	PCA	VA (V4)	BA
Group I	57,2±8,4	54,3±7,3	38,6±5,7	37,5±6,1	36,4±4,9
Group II	51,3±9,2	51,6±5,8	36,5±4,7	30,6±5,8	31,4±6,2
CG	62,6±10,1	52,3±6,7	36,5±5,7	34,7±9,1	38,9±8,4

**Table II.** Indicators of CVR in patients with CBI

	CrCO2	CrO2	IVMR	CrFNT
Group I	1,17±0,04*	0,34±0,04	62,9±7,5*	0,11±0,05
Group II	1,14±0,03*	0,26±0,03*	54,2±8,8*	0,05±0,03*
CG	1,28±0,06	0,36±0,03	81,5±6,9	0,16±0,04

\* p < 0,05 compared with CG

**Table III.** Indexes RIRP in patients with SDI, II and KA

	SDI	II	KA
Group I	1,27±0,03*	1,26±0,05*	1,20±0,04
Group II	1,32±0,05*	1,29±0,06*	1,19±0,03
CG	1,18±0,03	1,18±0,03	1,18±0,03

\*p < 0,05 compared with CG

autonomic support of activity (ASA) were characterized by excess (41.5%) or deficiency (47.1%). In the presence of subcompensated CBI, the latter increased up to 95.8% (p<0.01). A similar trend occurred in the presence of menopausal syndrome in women (59.4%, p < 0.05), while ASA was always pathological. Clinically significant depression was registered in 58.1% of cases, and 80.5% of them were women (p < 0.05), and in the presence of menopausal syndrome it reached 94.4% (p < 0.05). With vagotonia, these indicators were 100.0% in men and 59.5% in women. Anxiety was defined in 30.6% without gender differences, more often at the compensated phase of CBI. In case of dyshormonal changes in women, anxiety occurred much more often (72.9%), with a predominance of sympathicotonic background (51.8%). The parameters of hemodynamics in patients of Group I did not differ significantly from the data of CG, only the values of blood flow in the MCA slightly lowered. In patients of Group II, a moderate decrease in blood flow indicators was observed in MCA, VA and BA. Analogous indicators in ACA and PCA did not significantly differ from normative ones. These changes can be explained by the appearance of local stenosing processes in the main arteries affecting cerebral hemodynamics in patients with subcompensated CBI (Table I).

Indicators of RI in VA exceeded the normative values in the direction of angiospasm, the average RI in Group II was more often interpreted as dystonic, and in Group I, persistent cerebral angiospasm was manifested more frequently, especially against the background of sympathicotonia (up to 0.73±0.09 units, p < 0.05), in cases of epytonia these indicators decreased (0.53±0.08 units),

and in cases of vagotonia it reached the minimum values (0.46±0.08 units, p < 0.05). In patients of Groups I and II, a significant decrease in CrCO2 indicators was observed (Group I – 1.17±0.04, CG – 1.28±0.06, p<0.05; Group II – 1.14±0.03, CG – 1.28±0.06, p<0.05). CrO2 values in Group I did not change significantly, in Group II they significantly decreased (Group I – 0.34±0.04, CG – 0.36±0.03, Group II – 0.26±0.03, CG – 0.36±0.03, p<0.05). The values of IVMR being an integral indicator of stability of cerebral autoregulation showed sharp decrease in Groups I and II (Group I – 62.9±7.5, CG – 81.5±6.9, p<0.05; Group II – 54.2 ±8.8, CG – 81.5±6.9, p<0.05). These changes were mostly characteristic of the patients of Group II. Also, hyporeactivity to FNT (0.05±0.03, CG – 0.16±0.04, p<0.05) was significantly manifested in Group II, which is the most sensitive indicator of the vasodilation function disorder at various phases of CVP (Table II).

According to X-ray examination, signs of stair descent instability (SDI) in CRS C2-C6 were defined in 19.3% of patients of Group I, in 22.1% of patients of Group II. Isolated instability (II) in CRS C2-C3 was observed in 5.1% of patients of Group I and in 6.3% of patients of Group II, in CRS C3-C4 – in 17.4% of patients of Group I and in 21.7% of patients of Group II, in CRS C4-C5 – in 4.5% of patients of Group I and in 4.3 patients of Group II, in CRS C5-C6 – in 3.2% of patients of Group I and in 5.6% of patients of Group II. Signs of Kimmerli's anomaly (KA) were observed in 8.7% of patients of Group I and 5.1% of patients of Group II. In the clinical group of patients with SDI, pronounced hyperreactivity to functional loads with rotation to the right and left

was defined (Group I –  $1.27 \pm 0.03$ , CG –  $1.18 \pm 0.03$ ,  $p < 0.05$ ; Group II –  $1.32 \pm 0.05$ , CG –  $1.18 \pm 0.03$ ,  $p < 0.05$ ). In patients with II indicators of reactivity index to rotation probes (RIRP) were also significantly increased (Group I –  $1.26 \pm 0.05$ , CG –  $1.18 \pm 0.03$ ,  $p < 0.05$ ; Group II –  $1.29 \pm 0.06$ , CG –  $1.18 \pm 0.03$ ,  $p < 0.05$ ). In the group with KA, relative normal reactivity to the right and left rotation was observed (Group I –  $1.20 \pm 0.04$ , CG –  $1.18 \pm 0.03$ ; Group II –  $1.19 \pm 0.03$ , CG –  $1.18 \pm 0.03$ ) (Table III).

## DISCUSSION

Ischemic damage to the brain due to degenerative-dystrophic changes in the cerebral spine is often accompanied by VD. The presence of the latter correlates with the degree of brain damage. Vestibulo-ataxic disorders were higher compared to CG and increased along with the degree of brain damage. An important aspect of the development of VD is autonomic dysfunction against the background of pathological autonomic characteristics with predominant parasympathetic orientation of AT, especially in the case of insufficiency of autonomic reactivity (RA) and pathological ASA. Such changes significantly increased in the presence of subcompensation of CBI. The correlation between psychoemotional disorders and changes in autonomic characteristics with VD against the background of CBI with initial regularities depending on the degree of brain damage was defined. The progression of CBI is facilitated by coronavirus infection, which, in addition to neurotropic, leads to vascular damage, especially in the structures of the vestibular apparatus, which is sensitive to ischemia and manifested in autonomic, psychoemotional dysfunctions, and cognitive deficits. Our studies confirm the conclusions

that coronavirus infection affects the vestibular analyzer both in the acute and post COVID periods of the infection [20,21], with changes in functional characteristics of the autonomic system [22], affects cerebral hemodynamics [23]. A characteristic hemodynamic feature in groups with compensated and subcompensated CBI is the presence of reduced perfusion in BA and VA. Changes in cerebral vascular reactivity with a decrease in CrO<sub>2</sub> and CrFNT indicators were characteristic of the subcompensated phase of CBI. Hyperactivity to rotational functional loads in both clinical groups has a high correlation with the presence of stair descent and, to a lesser extent, isolated instability in CS.

## CONCLUSIONS

1. The occurrence of VD is facilitated by the presence of autonomic dysfunction and degenerative-dystrophic changes in the CS, especially in case of subcompensation of CBI.
2. Psychoemotional changes were a characteristic feature of patients with VD against the background of CBI and had certain regularities depending on the phase of CBI.
3. Suffered coronavirus infection contributes to the progression of VD and further decompensation of CBI due to direct damage to the autonomic and vascular systems of the brain.
4. Changes in cerebral hemodynamics in the form of reduced perfusion in BA and VA, a decrease in cerebrovascular reactivity, and an increase in reactivity to rotational functional load were determined in patients with VD against the background of subcompensated CBI.

## REFERENCES

1. Tong X., Yang Q., Ritchey M.D. et al. The Burden of Cerebrovascular Disease in the United States. *Prev Chronic Dis.* 2019;16:E52. doi: 10.5888/pcd16.180411.
2. Stoyanov O.M., Vastianov R.S., Kalashnykov V.Y. et al. Rol vehetatyvnoi systemy u formuvanni khronichnoi ishemii mozku, porushen tserbralnoi hemodynamiky, avtonomnomu rehuliuvani. [The role of the autonomic system in the formation of chronic brain ischemia, cerebral hemodynamic disorders, autonomic rehabilitation]. *Ukrainskyi visnyk psikhonevrolohi/ Ukrainian Herald of Psychoneurology.* 2022; 30 (112): 39-40. doi:10.36927/2079-0325-V30-is3-2022-33. (in Ukrainian).
3. Barpanda S. Pathophysiology and Epidemiology of Cerebrovascular Disease. *International Journal of Collaborative Research on Internal Medicine & Public Health Editorial.* 2021; 13 (7): 1-2.
4. Muratova T., Khramtsov D., Stoyanov A., Vorokhta Yu. Clinical epidemiology of ischemic stroke: global trends and regional differences. *Georgian Medical News.* 2020; 2 (299): 83-86. doi: 10.26693/jmbs05.01.149.
5. Bakumenko I.K., Son A.S., Stoyanov A.N., Vastyanov R.S. Vestibulyarnye disfunkcii i psihovegetativnye rasstrojstva pri ishemii mozga na fone shejnogo osteohondroza i ih patogeneticheski orijentirovannaya korrekciya. [Vestibular dysfunctions and psycho-vegetative disorders in brain ischemia on the background of cervical osteochondrosis and their pathogenetically oriented correction]. *Neurophysiology.* 2014; 46 (3): 300-303. (in Ukrainian).
6. Trinus K.F., Klaussen K. Mizhnarodnij klinichnij protokol z prisinkovih porushen (zapamorochen). [International clinical protocol of vestibular disorders (dizziness)]. *Shidno-yevropejskij nevrologichnij zhurnal/ East European Journal of Neurology.* 2015;4: 4-47 (in Ukrainian).

7. Dougherty J.M., Carney M., Hohman M.H., Emmady P.D. Vestibular Dysfunction. 2022 Aug 29. In: StatPearls. Treasure Island (FL): StatPearls Publishing. 2022.
8. Demydas O.V., Tkachenko O.V. Klinichni kharakterystyky stanu vehetatyvnoi nervovoi systemy ta yikh koreliatsiini vzaiemozv'iazky z psykometrychnymy pokaznykamy u patsientiv iz vyrazkoiu dvanadtsiatypaloi kyshky v stadii zahostrennia ta remisii [Clinical characteristics of the state of the autonomic nervous system and their correlation with psychometric indicators in patients with duodenal ulcer in the stage of exacerbation and remission]. Ukr Med Chasopys/ Ukrainian Medical Journal. 2022;4(150):1-4. (in Ukrainian). doi: 10.32471/umj.1680-3051.150.232312.
9. Ellul M.A., Benjamin L., Singh B. et al. Neurological associations of COVID-19. Lancet Neurol. 2020;19(9):767-783. doi: 10.1016/S1474-4422(20)30221-0.
10. Dries D.J., Hussein H.M. Coronavirus Disease 2019 and Stroke. Air Med J. 2021; 40(2):92-96. doi: 10.1016/j.amj.2020.12.003.
11. Fifi J.T., Mocco J. COVID-19 related stroke in young individuals. Lancet Neurol. 2020;19(9):713-715. doi: 10.1016/S1474-4422(20)30272-6.
12. Ostroumova T.M., Chernousov P.A., Kuznetsov I.V. Kognitivnye narusheniya u vyzhivshih posle COVID-19 [Cognitive impairment in COVID-19 survivors]. Nevrologiya, neiropsikhiatriya, psikhosomatika/ Neurology, Neuropsychiatry, Psychosomatics. 2021;13(1):126-130. doi: 10.14412/2074-2711-2021-1-126-130. (in Russian).
13. Obaid M., Flach C., Marshall I. et al. Long-Term Outcomes in Stroke Patients with Cognitive Impairment: A Population-Based Study. Geriatrics (Basel). 2020;5(2):32. doi: 10.3390/geriatrics5020032.
14. Stoyanov O.M., Son A.S., Vastyanov R.S. et al. Cognitive impairment restoration in patients suffered with stroke during the post-COVID period. Journal of Education, Health and Sport. 2022;12(2):336-343. doi: 10.12775/JEHS.2022.12.02.034.
15. Karaban I., Hasiuk T., Karasevych N. COVID-19 and Parkinson's disease. International Neurological Journal, 2022;18(3):46-50. doi:10.22141/2224-0713.18.3.2022.943.
16. Varga Z., Flammer A.J., Steiger P. et al. Endothelial cell infection and endotheliitis in COVID-19. Lancet. 2020;395(10234):1417-1418. doi: 10.1016/S0140-6736(20)30937-5.
17. Paniz-Mondolfi A., Bryce C., Grimes Z. et al. Central nervous system involvement by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2). J Med Virol. 2020; 92(7):699-702. doi: 10.1002/jmv.25915.
18. Sbornik metodik i testov dlya issledovaniya vegetativnogo otdela nervnoj sistemy (red. Yu.L. Kurako). [Collection of methods and tests for the study of the autonomic nervous system (Ed. Yu.L. Kurako)]. Odessa, OGMU. 1999, 192 p (in Russian).
19. Zigmond A.S., Snaith R.P. The hospital anxiety and depression scale. Acta Psychiatr Scand. 1983;67:361-370.
20. Potts R., Li H. Vestibular Dysfunctions Related to COVID-19 Infection: A Clinical Case Report (P9-9.003). Neurology. 2022; 98 (18): 2863.
21. Pazdro-Zastawny K., Dorobisz K., Misiak P. et al. Vestibular disorders in patients after COVID-19 infection. Front. Neurol. 2022; 13:956515. doi: 10.3389/fneur.2022.956515.
22. Dani M., Dirksen A., Taraborrelli P. et al. Autonomic dysfunction in 'long COVID': rationale, physiology and management strategies. Clin Med (Lond). 2021;21(1):e63-e67. doi: 10.7861/clinmed.2020-0896.
23. Sonkaya A.R., Öztrk B., Karadağ Ö. Cerebral hemodynamic alterations in patients with Covid-19. Turk J Med Sci. 2021;51(2):435-439. doi: 10.3906/sag-2006-203.

*Study of the peculiarities of the state of the central nervous system and central nervous system in vestibular dysfunctions in patients with chronic brain ischemia. State registration number: 0115U006651. Odesa National Medical University. Implementation period: 2017-2021.*

#### **ORCID and contributionship:**

Alexander N. Stoyanov: 0000-0002-3375-0452 <sup>A, C, D</sup>

Serhii S. Mashchenko: 0000-0001-6766-1231 <sup>A, C, D</sup>

Valeriy I. Kalashnikov: 0000-0002-7012-1698 <sup>A, B, D</sup>

Rooslan S. Vastyanov: 0000-0001-8585-2517 <sup>E, F</sup>

Alexander R. Pulyk: 0000-0002-8717-047X <sup>A, F</sup>

Tamara Andreeva: 0000-0002-4357-320X <sup>B, D</sup>

Olena O. Kolesnik: 0000-0003-0296-4447 <sup>C, E</sup>

#### **Conflict of interest:**

*The Authors declare no conflict of interest.*

---

## **CORRESPONDING AUTHOR**

**Valeriy I. Kalashnikov**

Kharkiv National Medical University

58 Amosova st., 61000 Kharkiv, Ukraine

tel: + 38-067-705-70-09

e-mail: dr.valkalash@gmail.com

**Received:** 27.08.2022

**Accepted:** 18.02.2023

---

**A** – Work concept and design, **B** – Data collection and analysis, **C** – Responsibility for statistical analysis, **D** – Writing the article, **E** – Critical review, **F** – Final approval of the article

 Article published on-line and available in open access are published under Creative Common Attribution-Non Commercial-No Derivatives 4.0 International (CC BY-NC-ND 4.0)