

# **GEORGIAN MEDICAL NEWS**

---

ISSN 1512-0112

NO 1(358) January 2025

---

TBILISI - NEW YORK



MONTHLY SCIENTIFIC JOURNAL

Georgian Medical News

საქართველოს სამედიცინო სიახლენი

## GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.  
Published since 1994. Distributed in NIS, EU and USA.

**GMN: Georgian Medical News** is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

**GMN: Медицинские новости Грузии** - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

**GMN: Georgian Medical News** – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

### WEBSITE

[www.geomednews.com](http://www.geomednews.com)

## К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html) В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

**При нарушении указанных правил статьи не рассматриваются.**

## REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: [http://www.nlm.nih.gov/bsd/uniform\\_requirements.html](http://www.nlm.nih.gov/bsd/uniform_requirements.html)  
[http://www.icmje.org/urm\\_full.pdf](http://www.icmje.org/urm_full.pdf)

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned  
Requirements are not Assigned to be Reviewed.**

## ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრამების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალებების შედეგების ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Larisa Melia, Revaz Sulukhia, Natia Jojua, Tinatin Gognadze, Nino Davidova. PRETERM BIRTH PREVENTION IN MULTIFETAL PREGNANCIES: A RETROSPECTIVE STUDY ON CERVICAL PESSARY EFFICACY.....	6-10
Ketevan Tsanava, Lali Khurtsia, Elene Shengelia, Gvantsa Qvariani, Luka Dangadze. DIAGNOSTIC CHALLENGE: COEXISTING MULTIPLE MYELOMA AND EXTRAMEDULLARY PLASMACYTOMA WITH RENAL AND HEPATIC INVOLVEMENT.....	11-14
Alghamdi Thamer, Khallufah Ahmed, Alghamdi Adel, Mohammed Al Shareef, Alzahrani Alaa, Alzahrani Faisal, Alghamdi Khader, Alghamdi Anmar. PREVALENCE, PATTERN, RISK FACTORS, AND MANAGEMENT OF ABDOMINAL AND INGUINAL HERNIAS IN KING FAHAD HOSPITAL AT AL-BAHA CITY, SAUDI ARABIA 2024.....	15-21
Samsonia M.D, Kandelaki M.A, Giorgadze T.A. TRANSMISSION OF RABIES VIRUS THROUGH A CONTACT LENS CONTAMINATED WITH SALIVA FROM AN INFECTED DOG (CASEREPORT).....	22-25
M.K. Osmnina, N.S. Podchernyaeva, V. A. Seraya, S.K. Kurbanova, O.V. Batureva, S.N. Chebusheva, O. V. Shpionkova, A.V. Polyanskaya, A.A. Skakodub, N.K. Ziskina. EFFICACY AND TOLERABILITY OF JANUS KINASE INHIBITOR TOFACITINIB IN JUVENILE LINEAR SCLERODERMA. CASE SERIES OF 5 PATIENTS.....	26-30
Huda Saif Al Dhaheri, Mohammad Fareed Khan. OCULAR MANIFESTATIONS IN A PATIENT WITH HIDRADENITIS SUPPURATIVA: A CASE STUDY.....	31-34
Hawar Sardar Hassan, Ahmed J. Allami, Duha Emad Taha, Hany Akeel Al-Hussaniy. BETTER DIAGNOSIS OF STROKE USING DIFFERENT B-VALUES IN MAGNETIC RESONANCE IMAGING.....	35-39
Tchernev G, Broshtilova V3, Kordeva S. INNOVATIONS IN DERMATOLOGIC SURGERY AND MELANOMA PATHOGENESIS: FROM THE PERSONALISED SURGERY TO THE CONCEPT OF GENOMIC MAPPING/ TARGETING VIA NITROSAMINES IN DRUGS: SPOTLIGHT ON CONTAMINATION OF ANGIOTENSIN CONVERTING ENZYME INHIBITORS (ACES) AND ANGIOTENSIN RECEPTOR BLOCKERS (ARBS).....	40-46
Yu.V. Boldyreva, I.A. Lebedev, E.V. Zakharchuk, E.A. Babakin, I.A. Aptekar. CONGENITAL HYPOTHYROIDISM: FROM THEORY TO PRACTICE- A CLINICAL CASE.....	47-49
Zana Lila, Sokol Krasniqi, Afrim Gjelij, Jacques Veronneau. COMPARATIVE ANALYSIS OF ENAMEL SURFACE WEAR INDUCED BY TWO CONCENTRATIONS OF ZIRCONIA PARTICLE TOOTHPASTE UNDER TWO ELECTRIC TOOTHBRUSHING MODALITIES.....	50-56
Rebecca Mills, Mohammad Zain Sohail, Hammad Sadique, Oliver Adebayo, Kanatheepan Shanmuganathan, Georgios Mamarelis, Shahanoor Ali, Ahmed Sanalla, Frank Acquaah, Abid Ali, Sadhin Subhash. VALID AND INFORMED CONSENT IN ORTHOPAEDIC SURGERY: A MULTICENTRE, REGIONAL SERVICE EVALUATION OF CURRENT UK PRACTICE.....	57-69
George Shaburishvili, Nikoloz Shaburishvili, Solomon Zeikidze. PROPORTION OF HEART FAILURE PATIENTS RECEIVING GUIDELINE RECOMMENDED DOSES OF BETA BLOCKERS IN GEORGIA: A STUDY ON TITRATION AND TOLERABILITY.....	70-77
Chaima Jemai, Haifa Zaibi, Tesnim Farhat, Nesrine Dhieb, Achwak Mehrez, Mouna Djebbi, Zohra Hadj Ali, Yosra Htira, Faika Ben Mami. STUDY OF THE ASSOCIATION BETWEEN ASTHMA, WEIGHT STATUS AND NUTRITIONAL INTAKE: RESULTS OF A TUNISIAN PILOTSURVEY.....	78-85
Robizon Tsiklauri, Tamar Jankhoteli, Maiko Chokheli, Ani Khachidze, Lela Kazarashvili, Nino Chkhaberidze, Ketevan Kavtaradze, Emzari Chachua, Mariam Vardoshvili. HEALTH RISK-FACTORS ASSOCIATED WITH LEAD EXPOSURE IN THE KVEMO KARTLI REGION OF GEORGIA.....	86-94
Najafbayli N.V. SEMANTICS AND DYNAMICS OF HEADACHE IN PATIENTS WITH CHIARI MALFORMATION TYPE I AFTER DECOMPRESSION SURGERY: EXPERIENCE FROM AZERBAIJAN.....	95-100
Hussamaldin Mohamed, Abdelmushin Abdelgadir, Ashraf Ismail, Osman Elsadig, Kiran Gopinath, Mosab Omer, Ayman Alfeel, Elryah. I. Ali, Mohamed M. Almaki, Ammar Abdelmola, Hussam Ali Osman, Huda Al-Obaidi, Abdelgadir Elamin Eltom, Marwan Ismail. EXPLORING THE ROLE OF C-REACTIVE PROTEIN IN PREECLAMPSIA AMONG HYPERTENSIVE PREGNANT WOMEN....	101-105
Tamar Shervashidze, Rusudan Kvanchakhadze, David abuladze, Liana Jashi, Miranda Shervashidze, Ilona Sakvarelidze, Manana Makharadze, Iamze Taboridze. THE IMPACT OF BARIATRIC SURGERY ON TYPE 2 DIABETES MELLITUS REMISSION IN THE GEORGIAN POPULATION.....	106-112

Wilfredo Chaviano-de la Paz, Dayani Arteaga-Guerra, Luis Enrique Remedios Carbonell, Raikel Fardales Rodriguez, Maidelis Prieto-Guerra, Michel Guillermo-Segredo, Maikel Santos-Medina, Geovedys Martinez-Garcia, Miguel Alejandro Rodríguez-Ramos. TEN-YEAR TRENDS IN REVASCULARIZATION, IN-HOSPITAL TREATMENTS, AND OUTCOMES IN PATIENTS WITH STEMI.....	113-120
Kubaevskaya D. M, Olennikov P. A, Ishmaev S. A, Balakireva E. V, Labazanov D. U, Boguslavets S. L, Beskadarov V. I, Zhidkov S. A, Budeykina I. N, Komolov D. A. FORMATION OF ARTIFICIAL BURNS IN WISTAR RATS TO EVALUATE THE EFFECTS OF DIFFERENT DRUGS.....	121-122
Tatiana V. Kirichenko, Irina Yu. Yudina, Maria V. Lukina, Tatiana B. Andrushchishina, Natalia V. Elizova, Alexander M. Markin, Yuliya V. Markina. IMMUNE RESPONSE OF CULTURED MONOCYTES OF ATHEROSCLEROTIC PATIENTS RECEIVING STATIN THERAPY.....	123-128
Yurko K.V, Chekhovska G.S, Gradil G.I, Katsapov D.V, Merkulova N.F, Mohylenets O.I, Bodnia I.P, Burma Ya.I, Tsyko O.V, Onikiienko O.L, Gargin V.V. DIAGNOSTIC MANAGEMENT OF PATIENTS WITH ONYCHOMYCOSES.....	129-133
Alyaa Abdulameer, Marwa Abdulzahra, Zainb Adel hashim. VARIATION OF ASTIGMATISM BETWEEN TEMPORAL AND SUPERIOR APPROACH IN PHACO SURGERY.....	134-137
Encarnación David Velásquez-Pasapera, Sofia Romero-Mederos, Jose Antonio Paredes-Arrascue. INTEROPERABILITY IN PERUVIAN BLOOD BANKS: PERCEPTION AND CHALLENGES FOR THE IMPLEMENTATION OF AN INTEGRATED INFORMATION SYSTEM.....	138-142
Tchernev G, Broshtilova V, Kordeva S. POLYPHARMACY, DRUG RELATED NITROSAMINE CONTAMINATION (BISOPROLOL/ PROPAFENONE) AND THE LINK TO LICHEN PLANUS/ SUBSEQUENT DEVELOPMENT OF KERATINOCYTE AND MUCOSAL CANCER/ ORAL LEUKOPLAKIA: PRESENTATION OF THE FIRST CASE AND UPDATE ON THE NEW PATHOGENETIC VISION.....	143-150
Ayhan Verit, Fatma Ferda Verit. “SCREAM” OF CYSTOLITHOTOMY IN HISTORY OF ART: PATIENT PERSPECTIVE.....	151-153
M.A. Rustamzade, N.M. Amiraliyev, K.N. Amiraliyev. EFFICIENT RECONSTRUCTION METHOD SELECTION IN LOWER LIP CANCER.....	154-157
Chaima Jemai, Radhouane Gharbi, Hajer Kandara, Ines Kammoun, Manel Jemel, Olfá Berriche, Faten Mahjoub, Henda Jamoussi. OBESITY AND THYROID FUNCTION IN OBESE WOMEN: A PILOT STUDY.....	158-162
Nazaryan R.S, Sosonna L.O, Iskorostenska O.V, Storozheva M.V, Fomenko Yu.V, Heranin S.I, Ohurtsov O.S, Nikonov A.Yu, Alekseeva V.V. ANATOMICAL FEATURES OF THE OSTIOMEATAL COMPLEX AND THEIR IMPACT ON COMPLICATIONS IN DENTAL IMPLANTATION.....	163-167

## ANATOMICAL FEATURES OF THE OSTIOMEATAL COMPLEX AND THEIR IMPACT ON COMPLICATIONS IN DENTAL IMPLANTATION

Nazaryan R.S<sup>1</sup>, Sosonna L.O<sup>1</sup>, Iskorostenska O.V<sup>2</sup>, Storozheva M.V<sup>1</sup>, Fomenko Yu.V<sup>1,2</sup>, Heranin S.P<sup>2</sup>, Ohurtsov O.S<sup>2</sup>, Nikonov A.Yu<sup>2</sup>, Alekseeva V.V<sup>1,3</sup>.

<sup>1</sup>Kharkiv National Medical University Kharkiv, Ukraine.

<sup>2</sup>V. Karazin Kharkiv National University Kharkiv, Ukraine.

<sup>3</sup>Kharkiv International Medical University, Kharkiv, Ukraine.

### Abstract.

The anatomical structure of the ostiomeatal complex (OMC) plays a crucial role in the success of dental implantation procedures, particularly in the maxillary region. This study examines the anatomical variations of the OMC and their impact on postoperative complications in dental implantation. A total of 400 spiral computed tomography (CT) scans of patients aged 18 to 95 years were analyzed to assess key anatomical structures, including the maxillary sinus, concha bullosa, uncinate processes, and sinus ostium dimensions. Findings revealed significant anatomical variability, with concha bullosa observed in 33.2% of cases, uncinate process variations in 31.6%, and a narrowed maxillary sinus ostium in 6.25% of cases. These structural differences were found to influence sinus ventilation and drainage, thereby increasing the risk of postoperative complications such as sinusitis and impaired osseointegration. Statistical analysis highlighted strong correlations between sinus volume, bone thickness, and OMC components, reinforcing the need for thorough preoperative assessment. The study underscores the importance of personalized surgical planning and advanced imaging techniques in mitigating complications and optimizing dental implantation outcomes.

**Key words.** Computer tomography, ostiomeatal complex, dental implantation, anatomical variability.

### Introduction.

The structure of the paranasal sinuses plays a crucial role in the success of surgical interventions in dentistry and maxillofacial surgery [1]. One of the major challenges in modern dentistry is the implantation of teeth, particularly in the maxillary region, due to the anatomical complexities of the maxillary sinus [2].

The proximity of the maxillary sinus floor to the alveolar ridge, variations in sinus morphology, and the presence of anatomical anomalies can significantly affect the success of dental implant placement [3]. In some cases, insufficient bone volume or sinus pneumatization necessitates additional surgical procedures, such as sinus lifting, to ensure implant stability [4].

A thorough understanding of the relationship between the maxillary sinus anatomy and potential complications in dental implantation is crucial for optimizing treatment planning and minimizing postoperative risks [5]. A particularly important factor post-surgery is the structure of the ostiomeatal complex (OMC), a key anatomical region of the skull responsible for the physiological ventilation of the paranasal sinuses [6]. Its structural characteristics significantly influence the likelihood of postoperative complications [7]. The contribution of each OMC component to sinus ventilation remains uncertain and unproven. Some studies have focused on the density of the

uncinate process in recurrent rhinosinusitis, but the question of this parameter in the absence of pathology remains unexplored.

During surgical treatments, the goal is usually to expand the area of the semilunar hiatus as much as possible to maximize sinus ventilation efficiency [8]. However, it is known that when the semilunar hiatus becomes too wide, turbulent airflow disappears, and sinus ventilation efficiency progressively decreases. Therefore, the question arises about determining the physiological dimensions of the semilunar hiatus and even identifying the critical upper limits that would still ensure adequate ventilation.

Given these considerations, the objective of our study was to examine the anatomical features of the ostiomeatal complex and their impact on complications in dental implantation.

### Materials and Methods.

This study was conducted based on the analysis of 400 spiral computed tomography (CT) scans of men and women aged 18 to 95 years. The participants were divided into groups according to the World Health Organization classification. None of the participants reported complaints related to the ear, nose, and throat (ENT) organs, and the CT scans were performed for reasons unrelated to ENT pathology (e.g., suspected stroke, which was later ruled out). Cases were excluded from the study if there was evidence of pathological changes in the paranasal sinuses (PNS), as well as in cases of childhood or pregnancy.

Data collection was conducted at the Kharkiv Research Institute of General and Emergency Surgery and the Merefá Central District Hospital, in accordance with a scientific and practical cooperation agreement (No. 173/10 18, dated October 18, 2018).

The study adhered to the ethical standards of the World Medical Association's Declaration of Helsinki on medical research involving human subjects. All participants were fully informed about the study and provided written informed consent. The research protocol was approved by the Bioethics Committee of Kharkiv National Medical University (Protocol No. 5, dated November 11, 2018).

**Imaging and Data Analysis:** Computed tomography was used not only to visualize the anatomical structures but also to analyze the spatial relationships between various elements of the PNS, the osteomeatal complex (OMC), and adjacent regions. Additionally, CT scans enabled measurements of bone thickness and density. Bone density was assessed using the Hounsfield scale [9], which allows modern CT devices to differentiate 4,096 shades of gray, representing different density levels in Hounsfield units (HU). According to this scale, water is assigned a value of 0 HU, while air is -1000 HU.

The study was conducted using a Toshiba Aquilion 64 [10] computed tomography scanner (Japan), a multi-slice CT scanner capable of simultaneously acquiring data from four 0.5 mm slices. This scanner is known for its high performance, featuring a full rotation time of up to 0.4 seconds. To ensure high-resolution imaging while minimizing radiation exposure, a 0,5 mm slice thickness was used for examining the paranasal sinuses. The device's advanced technology optimizes X-ray using, providing high-quality images with a reduced radiation dose.

For image analysis, the Radiant DICOM Viewer software (version 4.6.9, 64-bit) was used [11]. This software offers a user-friendly interface and high-performance capabilities for viewing medical images in DICOM PACS format.

### Statistical Analysis:

Statistical analysis was performed using methods of variational statistics. The normality of data distribution was assessed using the Shapiro-Wilk test. The prevalence of different anatomical variants of the PNS and OMC was calculated as a percentage of the total study population.

The range of individual variability in PNS and OMC structure was assessed using the following statistical parameters:

- M – average value.
- $\sigma$  – standard deviation.
- Cv – coefficient of variation.
- mCv – standard error of the coefficient of variation.
- Min – minimum value in the sample.
- Max – maximum value in the sample.

A correlation analysis was also conducted. The Pearson correlation coefficient ( $r$ ) was used to examine the relationship between Haller's cell dimensions and the width of the semilunar hiatus. The statistical significance of the correlation was determined using Student's t-test.

### Morphometric Analysis:

Several morphometric parameters were measured, including:

- The longitudinal and transverse dimensions of the middle turbinate.
- Spatial orientation and attachment level of the middle turbinate.
- Volume and dimensions of the uncinate process.
- Bone density of the uncinate process.
- Distance between the middle turbinate and the semilunar hiatus.

Additionally, the spatial arrangement of the semilunar hiatus and its potential connections with the sinuses were analyzed. These anatomical features are critical, as they may contribute to the chronicity of inflammatory conditions and influence the surgical approach in functional endoscopic sinus surgery.

### Results and Discussion.


Paradoxical deviation of the middle turbinate was observed in 19 patients (7.6%). A pneumatized middle turbinate (concha bullosa) was detected in 83 patients (33.2%), accounting for 3.25% of cases, resulting from the early development of an air cell within the anterior ethmoid labyrinth.

Among these, a multichambered concha bullosa (see Figure 1) was found in 14 individuals (5.6%).



**Figure 1.** CT. Coronal reconstruction. Bulla of the middle turbinate on both sides, on the left – two-chamber concha bullosa.

In 39 patients (15.6%), the concha bullosa was significantly enlarged, completely obstructing the middle nasal passage. It extended toward the uncinate process and the ethmoid bulla, effectively blocking the communication between the paranasal sinuses and the nasal cavity.



**Figure 2.** Pneumatization of the uncinate process on the left. SCT. Coronal reconstruction.

Borderline anatomical variants (see Figure 2) of the uncinate process were identified in 57 patients (31.6%) on CT scans. These included:

- Pneumatization in 18 patients (7.2%).
- Hyperplasia in 22 patients (8.8%).
- Paradoxical bending and displacement into the depth of the middle nasal meatus in 17 patients (6.8%).
- Hypoplasia 0,8%.

Asymmetry in the superior portions of the Uncinate processes was noted in 22.8% (57 patients), with variations in attachment sites. Agger nasi cells were hypertrophied in 9.6% (24 patients), always appearing as unilocular structures.


Haller cells were present in 1.75% (7 patients) and were associated with a narrower maxillary sinus ostium, increasing the risk of obstruction.

Narrowing of the natural ostium (<4 mm) was found in 6.25% (25 patients), which could impair sinus ventilation and drainage, thereby increasing the risk of postoperative complications.

The correlation between these parameters was also calculated. The highest degree of correlation was found between the thickness of the upper walls and the volume of the sinuses ( $r = 0.96$ ).


The dimensions of the middle turbinate are shown in Figure 3.

Dimensions of the uncinate process, distance from the gray turbinate to the semilunar foramen.



**Figure 3.** Dimensions of the middle turbinate in different age groups.

The physiological function of the paranasal sinuses, particularly the maxillary sinuses, is closely dependent on the size and efficiency of their natural openings (Figure 4), all of which drain into the middle nasal passage. In 0.5% of cases, the presence of an additional opening was observed, which can significantly disrupt normal maxillary sinus function.



**Figure 4.** CT of the paranasal sinuses. Coronal projection (the arrow indicates the semilunar hiatus).

The size of these openings plays a crucial role, ranging from 3 to 10 mm. Notably, 87% of subjects had a semilunar hiatus larger than 4 mm, with the most common size being 5 mm [12].

Our study revealed significant anatomical variability in the osteomeatal complex (OMC) and its potential impact on complications during dental implantation [13]. Anatomical variations such as paradoxical curvature of the middle turbinate, concha bullosa, hyperplasia and pneumatization of the uncinate process, as well as differences in the size of the natural ostium, can significantly influence nasal aerodynamics, paranasal sinus ventilation, and the functional state of the mucosa [14].

Concha bullosa was observed in 33.2% of cases, with larger formations contributing to obstruction of the middle nasal passage and impaired maxillary sinus drainage [15]. This obstruction may increase the risk of inflammatory complications following dental implantation, including chronic sinusitis and acute sinus infections. Additionally, hyperplasia or pneumatization of the uncinate process, detected in 16% of participants, can further disrupt normal OMC function. This is supported by a significant negative correlation (-0.59) between the size of the uncinate process and the width of the semilunar hiatus.

One of the most critical factors influencing complication risk is the individual anatomical variability of the semilunar hiatus width [16]. In 6.25% of cases, this width was less than 4 mm—a threshold considered critical for proper sinus drainage [17]. Patients with a narrower semilunar hiatus are at a higher risk of inflammatory complications post-implantation. Furthermore, Haller cells were identified in 1.75% of cases, correlating with a reduction in the natural maxillary sinus ostium size, potentially leading to sinus obstruction [18].

Our findings also demonstrated a strong correlation between paranasal sinus volume, sinus wall thickness, and OMC parameters [19]. Notably, a high correlation ( $r = 0.96$ ) between sinus volume and upper wall thickness highlights the potential risks associated with sinus lift procedures. This emphasizes the necessity of preoperative bone density assessment before implantation [20,21].

One of important conclusion of described work is one more emphasis about common work of different specialist (dentist, radiologist, morphologist) even in period of wide digital medicine using [22-25] with start of implementation of artificial intelligence.

### Clinical Implications.

From a clinical perspective, our findings suggest several key considerations:

- 1. Preoperative Evaluation:** Patients with pronounced anatomical variations (e.g., uncinate process hyperplasia, concha bullosa, or a narrow semilunar hiatus) require thorough pre-implantation assessment.
- 2. Comprehensive Planning:** Preoperative planning should extend beyond bone volume analysis to include an evaluation of OMC ventilation and drainage function.
- 3. Advanced Imaging:** The use of spiral computed tomography (SCT) should be a standard diagnostic tool for identifying anatomical variations and determining optimal surgical strategies.

### Conclusion.

Understanding the anatomical features of the ostiomeatal complex is essential for predicting and minimizing complications

**Table 1.** Anatomical variability of the components of the osteomeatal complex in people of different ages and sexes.

Indicator	More than 90 years old		75-89 years		60-74 years		45-59 years		18-44 years	
	Male	Female	Male.	Female.	Male	Female	Male.	Female	Male.	Female
Volume of the uncinatate process, x10 <sup>-3</sup> m <sup>3</sup>	11940±810	9533±110	9870±930	8331±976	12468±1125	11876±639	13113±1125	13111±1212	14890±979	15111±2311
Density of the uncinatate germ, Hu	139±21	115±21	147±18,9	101±71	199±25	116±12	154±57	160±71	183±22	192±12
Longitudinal size of the middle turbinatate, x10 <sup>-3</sup> m	16,4 ±2,9	13,1±1,99	13,5±2,1	12,07±2,2	10,1±3	14,01±2,3	10,8±2,98	11,022±3,04	12,11±2,2	11,9±3,1
Transverse size of the middle turbinatate x10 <sup>-3</sup> m	5,47±1,2	3,65±1,87	2,88±0,61	4,9±1,2	4,96±0,88	4,12±0,29	4,41±2,02	3,346±0,79	6,08±0,99	4,33±0,93
Dimensions of the hiatus semilunaris x10 <sup>-3</sup> m	4,99±1,01	4,41±0,9871	4,56±1,06	5,01±0,77	4,54±0,8	3,99±0,97	5,14±1,4	6,02±1,8	7,6±1,28	4,54±0,5

in dental implantation. Patients with pronounced anatomical variations, such as hyperplasia of the Uncinate process, concha bullosa, or a narrow sinus ostium, require more thorough preoperative evaluation. Treatment planning should incorporate both bone volume analysis and an assessment of the ventilation and drainage functions of the ostiomeatal complex.

Future research should focus on refining imaging techniques and developing tailored surgical approaches to mitigate the risks associated with anatomical variations in the ostiomeatal complex.

**Funding:** This research received no external funding

**Conflict of interest statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**REFERENCES**

1. Bhalla N, Dym H. Update on Maxillary Sinus Augmentation. Dent Clin North Am. 2021;65:197-210.
2. Danesh-Sani SA, Loomer PM, Wallace SS. A comprehensive clinical review of maxillary sinus floor elevation: anatomy, techniques, biomaterials and complications. Br J Oral Maxillofac Surg. 2016;54:724-730.

3. Aldelaimi TN, Khalil AA. Maxillary Sinus Augmentation. *J Craniofac Surg*. 2016;27:e557-9.
4. Cruz RS, Lemos CAA, Batista VES, et al. Short implants versus longer implants with maxillary sinus lift. A systematic review and meta-analysis. *Braz Oral Res*. 2018;32:e86.
5. Testori T, Weinstein T, Taschieri S, et al. Risk factors in lateral window sinus elevation surgery. *Periodontol* 2000. 2019;81:91-123.
6. Alekseeva VV, Nechiporenko AS, Lupyr AV, et al. A method of complex evaluation of morphological structure of ostiomeatal complex components, lower wall of maxillary and frontal sinuses. *wiad lek*. 2020;73:2576-2580.
7. Whyte A, Boeddinghaus R. Imaging of odontogenic sinusitis. *Clin Radiol*. 2019;74:503-516.
8. von Arx T, Lozanoff S, Bornstein MM. Extraoral anatomy in CBCT – a literature review. Part 1: Nasoethmoidal region. *Swiss Dent J*. 2019;129:804-815.
9. DenOtter TD, Schubert J. Hounsfield Unit. 2023 Mar 6. In: *StatPearls* [Internet]. Treasure Island (FL): StatPearls Publishing; 2025.
10. Bauknecht HC, Jach C, Bohner G, et al. Impact of image noise levels, scout scan dose and lens shield on image quality and radiation exposure in z-axis dose-modulated neck MSCT on 16- and 64-slice Toshiba Aquilion scanners. *Eur J Radiol*. 2010;73:439-43.
11. Kats L, Goldman Y, Kahn A. Automatic detection of image sharpening in maxillofacial radiology. *BMC Oral Health*. 2021;21:411.
12. Hacking C, Jones J, St-Amant M, et al. Ostiomeatal complex narrowing due to anatomical variations. *Radiopaedia.org*. 2025.
13. Mirhadi S, Khojastepour L, Mirhadi S, et al. Anatomical variations of ostiomeatal complex in CBCT of patients seeking rhinoplasty. *J Dent (Shiraz)*. 2015;16:42-48.
14. Das S, Bhutia O, Saha R, et al. Osteomeatal complex: A study of its anatomical variation among patients attending North Bengal Medical College and Hospital. *J Clin Diagn Res*. 2015;9:AC04-7.
15. Prasanna LC, Mamatha H. Variations of ostiomeatal complex and its applied anatomy: a CT scan study. *Indian J Sci Technol*. 2010;3:904-907.
16. L Sandhu R, Kheur MG, Lakha TA, et al. Anatomic variations of the osteomeatal complex and its relationship to patency of the maxillary ostium: A retrospective evaluation of cone-beam computed tomography and its implications for sinus augmentation. *J Indian Prosthodont Soc*. 2020;20:371-377.
17. Riello A, Boasquevisque E. Anatomical variants of the ostiomeatal complex: tomographic findings in 200 patients. *Radiol Bras*. 2008;41:149-154.
18. Güngör G, Okur N, Okur E. Uncinate Process Variations and Their Relationship with Ostiomeatal Complex: A Pictorial Essay of Multidetector Computed Tomography (MDCT) Findings. *Pol J Radiol*. 2016;81:173-180.
19. Chmielewski PP. Clinical anatomy of the paranasal sinuses and its terminology. *Anat Sci Int*. 2024;99:454-460.
20. El-Din WAN, Madani GA, Abdel Fattah IO, et al. Prevalence of the anatomical variations of concha bullosa and its relation with sinusitis among Saudi population: a computed tomography scan study. *Anat Cell Biol*. 2021;54:193-201.
21. Lyndin M, Gluschenko N, Sikora V, et al. Morphofunctional features of articular cartilage structure. *Folia Med Cracov*. 2019;59:81-93.
22. Alekseeva V, Nechiporenko A, Frohme M, et al. Intelligent Decision Support System for Differential Diagnosis of Chronic Odontogenic Rhinosinusitis Based on U-Net Segmentation. *Electronics (Switzerland)*. 2023;12.
23. Nechiporenko A, Frohme M, Strelchuk Y, et al. Galvanic Skin Response and Photoplethysmography for Stress Recognition Using Machine Learning and Wearable Sensors. *Applied Sciences*. 2024;14:11997.
24. Gargin V, Radutny R, Titova G, et al. Application of the computer vision system for evaluation of pathomorphological images. 2020 IEEE 40th International Conference on Electronics and Nanotechnology, ELNANO 2020 – Proceedings. 2020:469-473.
25. Nechiporenko AS, Nazaryan RS, Semko GO, et al. Application of spiral computed tomography for determination of the minimal bone density variability of the maxillary sinus walls in chronic odontogenic and rhinogenic sinusitis. *Ukrainian journal of radiology and oncology*. 2021;29:65-75.