

• • • • •

• • • • •

•

• • • • •

•

• • • • •

• • • • •

• • • • •

•

•

• • • • •

•

•

• • • • •

• • • • •

•

• • • • •

()

•

, , .
:
-

[1].

-

.
-
,
,
,
:
36% (« »,)
,) « 1» (-
« »,).

+ , - ,

Staphylococcus spp., Streptococcus spp., Proteus vulgaris, Candida albicans,

, , .
.
,
, , , , .
, , , .
, ,

[2].

II – III

(- OHI-S (J.C. Green, J.R. Vermillion, 1964);
 - PI (A. L. Russel, 1956), (Papilla Bleeding
 Index, Mühlemann – Saxer, 1975), (Kötzschke, 1968),
) 18
 2 : 1 – (12) 2-
 (6).

1-2 « »,

« 1». 2

0,05%.

1.

1

	OHI-S	PI	PBI	
	2,2±0,3	4,9±0,15	3,00±0,03	4,6±0,3 6,7±0,4

II – III .

,
 ,
 ,
 - () .
 :
 , 2/3

II – III

, (, 1994).

1

: , ,
 ,

2

5- 10-

2.

2

-	OHI-S		PI		PBI			
	5	10	5	10	5	10	5	10
		1,8±0,3	1,7±0,1	2,5±0,26	2,3±0,29	1,5±0,03	1,4±0,12	2,4±0,13 3,5±0,18
	2,0±0,35	1,9±0,4	3,1±0,15	2,9±0,26	2,8±0,9	2,6±0,35	4,4±0,41 6,2±0,13	4,3±0,05 5,9±0,07

« »

« »

(-) 3-4 1

2

: ()
) ,

1

OHI-S, PI, PBI (.

. 2)

(< 0,05).

1-9



. 1. ., 49 .



. 2.



. 3. 36%



. 4.



. 5.



. 6.



.7.

36%

1. - : ‘ , 2000.-
464 .
2. . . .
/ //
// 4-
(27-28 2012),- : « »,-
2012.- .69-71.

• ” • ” • •

36%

,

.

:

,

,

• ”

• ”

• •

36%

.

:

,

,

Sokolova I. I., Denysova, Ye.G., Stoyan Ye.U.

COMBINED CONSERVATIVE TREATMENT OF PROLIFERATIVE PERIODONTAL
DISEASES

Kharkov national medical university

The effectiveness was studied of combined use of 36% solution polikrezulena and a polyvalent drug based haemostatic sponge in the topical treatment of generalized periodontitis, in the acute stage during the diagnosis of somatic diseases and microbiology of periodontal pockets.

Keywords: periodontitis, polikrezulen, haemostatic sponge