



ISSN 2226-1230 (Print)
ISSN 2413-4260 (Online)

89773

НЕОНАТОЛОГІЯ, ХІРУРГІЯ ТА ПЕРИНАТАЛЬНА МЕДИЦИНА

NEONATOLOGY, SURGERY AND PERINATAL MEDICINE

T.XIV, №2(52), 2024

DOI: 10.24061/2413-4260.XIV.2.52.2024

**АКТУАЛЬНІ ПИТАННЯ ОРГАНІЗАЦІЇ МЕДИЧНОЇ ДОПОМОГИ
В УКРАЇНІ. ДОДИПЛОМНА ТА ПІСЛЯДИПЛОМНА
МЕДИЧНА ОСВІТА**

- Ю. Г. Антупкін, Т. К. Знаменська, Р. В. Марушко,
О. О. Дудіна, О. В. Воробйова, Н. Ю. Бондаренко*
НАСЛІДКИ ВПЛИВУ НЕГАТИВНИХ ФАКТОРІВ ВОЄННОЇ
АГРЕСІЇ НА МЕДИЧНЕ ЗАБЕЗПЕЧЕННЯ ТА ЗДОРОВ'Я
НОВОНАРОДЖЕНИХ ПІД ЧАС ВІЙНИ 5
- О. Ковальова, С. Дудник, В. Похилько, Ю. Чернявська,
С. Цвіренко, А. Давиденко*
ПЕРИНАТАЛЬНА АСПІКСІЯ ТА ЇЇ НАСЛІДКИ: ВИКЛИКИ
СЬОГОДЕННЯ В УКРАЇНІ..... 12
- О. С. Полянська*
ФІЗИЧНА ТА РЕАБІЛІТАЦІЙНА МЕДИЦИНА В УКРАЇНІ:
РОЗВИТОК ТА ДОСЯГНЕННЯ 19
- Л. І. Вакулєнко, Л. П. Бадогіна, О. Ю. Оболонська,
А. В. Різник, С. В. Самсоненко*
ІНОВАЦІЙНІ МЕТОДИ У ВИКЛАДАННІ ПЕДІАТРІЇ
ЗДОБУВАЧАМ ВИЩОЇ МЕДИЧНОЇ ОСВІТИ
НА V ТА VI КУРСАХ: ДІАЛОГОВЕ НАВЧАННЯ..... 23

**РЕЗУЛЬТАТИ ДИСЕРТАЦІЙНИХ
ТА НАУКОВО-ДОСЛІДНИХ РОБІТ**

НЕОНАТОЛОГІЯ, ПЕДІАТРІЯ

- О. С. Годованець, Ю. М. Нечитайло*
КЛІНІЧНА ХАРАКТЕРИСТИКА ТА МОЖЛИВОСТІ
ЛАБОРАТОРНОЇ ДІАГНОСТИКИ ГАСТРОІНТЕСТИНАЛЬНИХ
ПОРУШЕНЬ ПРИ ПЕРИНАТАЛЬНІЙ ПАТОЛОГІЇ
У ПЕРЕДЧАСНО НАРОДЖЕНИХ ДІТЕЙ 28
- О. С. Моштук, М. М. Кісельова*
ОСОБЛИВОСТІ МІКРОБНОЇ КОЛОНІЗАЦІЇ КИШЕЧНИКУ
ПРИ КОЛЬКАХ У НЕМОВЛЯТ, В АНАМНЕЗІ ЯКИХ БУЛА
АНТИБІОТИКОТЕРАПІЯ..... 34

ДИТЯЧА ХІРУРГІЯ

- О. М. Горбатюк, Д. С. Солейко, К. М. Шатрова, Н. П. Солейко*
КОМПЛЕКСНЕ ЛІКУВАННЯ ПЕРИАНАЛЬНОЇ ХВОРОБИ
КРОНА У ДІТЕЙ 42

СТОМАТОЛОГІЯ

- В. В. Гороховський, О. В. Деньга, С. В. Скульська, В. М. Почтар*
РЕГІОНАЛЬНІ ОСОБЛИВОСТІ ТЕРМІНІВ ПРОРІЗУВАННЯ
ПОСТІЙНИХ ЗУБІВ У РАНЬОМУ ЗМІННОМУ ПРИКУСІ
В ДІТЕЙ МІСТА ОДЕСИ..... 50

АКУШЕРСТВО ТА ГІНЕКОЛОГІЯ

- В. Ліхачов, О. Тарановська, І. Жабченко, В. Оксютя,
В. Палапа, Е. Крутківа*
ПЕРЕДУМОВИ САМОВІЛЬНОГО ВИКИДНЯ В РАННІХ
ТЕРМІНАХ ВАГІТНОСТІ, ЩО НАСТАЛАЛА НА ФОНІ
ХРОНІЧНОГО ЕНДОМЕТРИТУ 57
- І. Лецишин, Л. Маркулан, П. Бик, Я. Сусак, Н. Мартинюк,
О. Охоцька*
ЯКІСТЬ ЖИТТЯ У ВАГІТНИХ З ХРОНІЧНИМИ
ПОВІЛЬНО-ТРАНЗИТОРНИМИ ЗАКРЕПАМИ 63
- І. Д. Кудін, В. В. Лазуренко, В. Ю. Прокопюк,
О. Ю. Железняков, М. Л. Панасовський*
ЕРИПТОЗ ЯК ЗВ'ЯЗОК МІЖ БАКТЕРІАЛЬНОЮ
ІНФЕКЦІЄЮ РЕПРОДУКТИВНОГО ТРАКТА
ТА АНЕМІЄЮ ВАГІТНИХ 70
- Б. В. Соколов, А. М. Бербець, О. М. Юзько*
ЯКІСТЬ ЖИТТЯ ТА РІВНІ МЕЛАТОНІНУ ТА СТЕРОЇДНИХ
ГОРМОНІВ В ПЛАЗМІ КРОВІ У ЖІНОК З ЛЕЙОМІОМОЮ
МАТКИ 78
- Н. В. Коцабин, А. В. Бойчук, Л. Б. Николин, Якимчук Ю. Б.*
АНАЛІЗ ВІЯВЛЕННЯ АНОМАЛЬНИХ
ГІСТЕРОСКОПІЧНИХ ЗНАХІДОК У ЖІНОК З НЕПЛІДДАМ. 86

**TOPICAL ISSUES OF MEDICAL CARE ORGANIZATION
IN UKRAINE. UNDERGRADUATE AND POSTGRADUATE
MEDICAL EDUCATION**

- Y. Antipkin, T. Znamenska, R. Marushko, O. Dudina, O. Vorobiova,
N. Bondarenko*
CONSEQUENCES OF THE INFLUENCE OF THE NEGATIVE
FACTORS OF MILITARY AGGRESSION ON THE MEDICAL
CARE AND HEALTH OF NEWBORN IN THE CONDITIONS
DURING THE WAR 5
- O. Kovalova, S. Dudnyk, V. Pokhilko, Yu. Chernyavska,
S. Tsvirenko, A. Davydenko*
PERINATAL ASPHYXIA AND ITS CONSEQUENCES:
TODAY'S CHALLENGES IN UKRAINE..... 12
- O. Polianska*
DEVELOPMENT OF PHYSICAL AND REHABILITATION
MEDICINE IN UKRAINE DURING THE PERIOD
OF MARTIAL STATUS 19
- L. Vakulenko, L. Badogina, O. Obolonska, A. Riznyk,
S. Samsonenko*
INNOVATIVE METHODS IN TEACHING PEDIATRICS
TO STUDENTS OF HIGHER MEDICAL EDUCATION
IN V AND VI YEARS: DIALOGUE TEACHING..... 23

**DISSERTATION RESULTS
AND RESEARCH WORKS**

NEONATOLOGY, PEDIATRIC

- O. Godovanets, Yu. Nechytailo*
CLINICAL CHARACTERISTICS AND POSSIBILITIES OF
LABORATORY DIAGNOSTICS OF GASTROINTESTINAL
DISEASES IN PERINATAL PATHOLOGY OF PREMATURE
INFANTS..... 28
- O. Moshtruk, M. Kiselyova*
FEATURES OF MICROBIAL COLONIZATION IN THE
INTESTINES IN COLIC IN INFANTS WITH A HISTORY OF
ANTIBIOTIC THERAPY 34

PEDIATRIC SURGERY

- O. Gorbatyuk, D. Soleiko, K. Shatrova, N. Soleiko*
COMPLEX TREATMENT OF PERINATAL CROHN'S
DISEASE IN CHILDREN 42

DENTISTRY

- V. Horokhovskiy, O. Dienha, S. Skulska, V. Pochtar*
REGIONAL FEATURES OF THE TERMS OF PERMANENT
TEETH ERUPTION IN EARLY MIXED DENTITION AMONG
CHILDREN OF ODESA..... 50

OBSTETRICS AND GYNECOLOGY

- V. Likhachov, O. Taranovska, I. Zhabchenko, V. Oksiuta,
V. Palapa, E. Krutikova*
PREREQUISITES FOR EARLY PREGNANCY LOSS
IN WOMEN WITH CHRONIC ENDOMETRITIS..... 57
- I. Leschyshyn, L. Markulan, P. Byk, Y. Susak, N. Martyniuk,
O. Okhotska*
QUALITY OF LIFE IN PREGNANT WOMEN WITH
CHRONIC TRANSIT CONSTIPATION 63
- I. Kudin, V. Lazurenko, V. Prokopiuk, O. Zhelezniakov,
M. Panasovskiy*
ERYPTOSIS AS A LINK BETWEEN BACTERIAL
REPRODUCTIVE TRACT INFECTION AND ANEMIA IN
PREGNANT WOMEN..... 70
- B. Sokolov, A. Berbets, O. Yuzko*
QUALITY OF LIFE AND PLASMA LEVELS OF MELATONIN
AND STEROID HORMONES IN WOMEN WITH UTERINE
LEIOMYOMA..... 78
- N. Kotsabyn, A. Boychuk, L. Nykolyn, Y. Yakymchuk*
ANALYSIS OF THE DETECTION OF ABNORMAL
HYSTEROSCOPIC FINDINGS IN WOMEN WITH INFERTILITY... 86

UDC: 618.3-022.7/9-06:616.155.194]-078
DOI: 10.24061/2413-4260. XIV.2.52.2024.11

ERYPTOSIS AS A LINK BETWEEN BACTERIAL REPRODUCTIVE TRACT INFECTION AND ANEMIA IN PREGNANT WOMEN

**I. Kudin, V. Lazurenko, V. Prokopiuk,
O. Zhelezniakov, M. Panasovskyi**

Kharkiv National Medical University
(Kharkiv, Ukraine)

Summary

Genital tract infections of various etiologies and anemia remain common in pregnant women. At present, there is a paucity of data on the impact of reproductive tract infections on the development of anemia in pregnant women, with particular emphasis on the contribution of eryptosis.

The aim of the current study was to assess the ability of bacterial, viral and fungal reproductive tract infections to trigger eryptosis of circulating erythrocytes in pregnant women.

Methods. The study included 47 pregnant women. According to the pathology of the infection, the patients were divided into three groups: 14 women with bacterial, 12 women with viral, 11 women with fungal infections of the reproductive tract and 10 pregnant women without extragenital and obstetric pathology (control subjects). The following methods were used to determine infection: bacteriologic, microscopic, polymerase chain reaction, immunoenzymatic, etc. Flow cytometry-based detection of eryptosis parameters (cell morphology, membrane phospholipid scrambling, and oxidative stress) was performed in pregnant women with reproductive tract infections of bacterial, viral, or fungal origin and in healthy individuals. Annexin V-FITC staining and 2',7'-dichlorodihydrofluorescein diacetate (H2DCFDA) staining were used to quantify phosphatidylserine exposure and reactive oxygen species (ROS) generation, respectively. Fluorescence was acquired using a BD FACSCanto™ II flow cytometer.

ANOVA test was used to compare numerical data of eryptosis assays. Post-hoc Bonferroni test was applied afterwards. Data were processed by Graph Pad Prism 5.0 software (USA). The difference was considered statistically significant at $p < 0.05$.

The research was conducted in accordance with the fundamental principles of Good Clinical Practice (GCP, 1996), the Council of Europe Convention on Human Rights and Biomedicine (April 4, 1997), the Helsinki Declaration of the World Medical Association on Ethical Principles for Medical Research Involving Human Subjects (1964-2008), as well as the Ministry of Health of Ukraine Order No. 690 dated September 23, 2009 (amended by the Ministry of Health of Ukraine Order No. 523 dated July 12, 2012).

The research design was discussed and approved at the meeting of the Medical and Ethical Committee of the Kharkiv National Medical University (Protocol No. 19 dated May 19, 2024).

The paper is an excerpt from the scientific research project of the Department of Obstetrics and Gynecology No. 2 at Kharkiv National Medical University, entitled «Optimisation of diagnostics, treatment of diseases of the reproductive system and pregnancy complications in women with extragenital pathology» (state registration number 0121U110923.)

Results. Erythropoiesis parameters were evaluated in pregnant women with bacterial, viral and fungal reproductive tract infections. Pregnancy-associated bacterial reproductive tract infections were associated with increased phosphatidylserine externalization in circulating erythrocytes, without ROS accumulation or cell shrinkage. Erythrocytes from patients with genital tract infections of viral or fungal origin show no signs of eryptosis, such as cell membrane scrambling, oxidative stress, or changes in cell morphology.

Conclusions. Reproductive tract infections of bacterial origin in pregnant women are associated with accelerated eryptosis, which may contribute to anemia in these women. Exclusively bacterial genital tract infection was shown to induce lipid membrane scrambling and hence eryptosis in erythrocytes from pregnant women. Phosphatidylserine externalization did not differ significantly between women with viral or fungal infections and healthy controls. Eryptosis induced by bacterial genital tract infection was not associated with cell shrinkage or ROS overproduction.

Key words: Reproductive Tract Infections; Erythrocyte; Anemia; Pregnancy; Eryptosis.

Introduction

Genital tract infections remain common in pregnant women, affecting up to 70 % of pregnancies, with widely varying prevalence rates reported in different studies [1-3]. Etiologically, lower tract infections can be caused by a variety of microorganisms [4]. Bacterial, viral, and fungal lower tract infections have been widely reported [5]. There is increasing evidence that candidiasis remains the most common genital tract infection in pregnant women [2, 6]. Lower tract infections have been associated with fetal infections and adverse pregnancy outcomes, including spontaneous abortion, preterm delivery, and perinatal complications [7, 8]. Thus, infections at the maternal-fetal interface remain of great clinical importance, and further

investigation of the molecular mechanisms and interplay governing host-pathogen interactions in genital tract infections in pregnant women should be encouraged [9]. In particular, anemia is considered to be common in pregnancy, being observed in up to 25 % of pregnancies [10, 11]. The relationship between anemia and UTIs in pregnancy has not been well studied. However, it has been reported that iron deficiency anemia is associated with susceptibility to UTI [12]. In this study, we hypothesized that eryptosis, a suicidal cell death mode of erythrocytes [13-15], may serve as a link between reproductive tract infections and anemia in pregnant women. Eryptosis has been shown to be a novel factor in the development of anemia in a variety of pathologies. In addition, it has recently been reported

to be a driver of anemia in chronic kidney disease [16]. In general, eryptosis is a cell death of mature erythrocytes that is morphologically similar to apoptosis but different from a signaling point of view [14, 17]. Eryptosis signaling is governed by intracellular Ca^{2+} elevation resulting in dysregulation of flippases and scramblases with subsequent phosphatidylserine externalization, activation of calpain resulting in cytoskeleton degradation and morphological changes in erythrocytes like cell shrinkage [13, 18]. Of note, ceramide and reactive oxygen stress (ROS) are considered important second messengers in eryptosis, which also contribute to the phospholipid membrane scrambling that is a hallmark of eryptosis [19-21]. It should be noted that phosphatidylserine externalization, an event that culminates eryptosis signaling, mediates the clearance of erythrocytic red blood cells from the circulation. This process, termed efferocytosis, which implies engulfment of erythrocytes by phosphatidylserine-associated mechanisms, underlies the pathogenesis of anemia in diseases associated with accelerated eryptosis.

The aim of the current study was to detect the impact of pregnancy-related genital tract infections in a causative agent-dependent manner on eryptosis of circulating mature erythrocytes, which has been hypothesized to contribute to anemia in pregnancy.

Materials and methods

Characteristics of patients.

The study included 47 pregnant women. According to the pathology of the infection, the patients were divided into three groups: 14 women (29.8 %) with bacterial, 12 women (25.5 %) with viral, 11 women (23.4 %) with fungal infections of the reproductive tract and 10 pregnant women (21.3 %) without extragenital and obstetric pathology (controls). There were no differences between the groups with regard to pregestational body mass index, age, parity, gestational weight gain and neonatal outcomes. The age of the studied pregnant women ranged from 20 to 37 years, the mean age of pregnant women with bacterial infection was 33.5 ± 4.1 years, pregnant women with viral infection – 28.2 ± 3.1 years, pregnant women with fungal infection – 24.7 ± 2.5 years, pregnant women in the control group – 25.1 ± 3.7 years. The following methods were used to determine infection: bacteriologic, microscopic, polymerase chain reaction, immunoenzymatic, etc. The diagnostics of infection agents was based on the result of the bacterioscopic research of the contents of the vagina, canal of the cervix, the urethra, bacteriological studies of vaginal secretion and identification of pathogens in the smears from the mucosa of the cervical canal by the method of polymerase chain reaction with the determination of the quantity infections agents.

Sample preparation.

Blood samples were collected in EDTA Vacutainers from pregnant women with genital tract infections and healthy volunteers. The samples were immediately used for flow cytometric detection of eryptosis markers. Briefly, samples were centrifuged and washed twice with phosphate-buffered saline (PBS; pH 7.4; BD, USA). Once red blood cell (RBC) suspensions were obtained,

a total of 2 μL of erythrocyte mass from each sample was resuspended in either 100 μL of PBS or annexin-binding buffer (Becton Dickinson) or PBS, depending on the staining procedure.

Membrane phospholipid scrambling detection.

RBCs resuspended in Annexin Binding Buffer as described above were stained with Annexin V-FITC (BD Pharmingen™ FITC Annexin V) for 15 minutes. After incubation, cells were resuspended in Annexin Binding Buffer. Fluorescence was acquired using a BD FACS Canto™ II flow cytometer (Becton Dickinson, USA): excitation wavelength was 488 nm; emission wavelength was 525 nm. To compare phospholipid scrambling, the percentage of cells displaying phosphatidylserine was analyzed [22-25].

ROS detection in erythrocytes.

Cells resuspended in PBS were stained with 5 μM 2',7'-dichlorodihydrofluorescein diacetate (H2DCFDA) provided by Invitrogen™ (USA). After incubation for 30 minutes, fluorescence was detected. The excitation and emission wavelengths were 488 nm and 525 nm, respectively. The mean fluorescence intensity of the ROS level-reflecting probe was analyzed to quantify ROS signaling [23-25].

Cell shrinkage analysis.

Forward scatter analysis was used to analyze cell morphology of erythrocytes from pregnant women with reproductive tract infections of bacterial, viral, or fungal origin. Since cell shrinkage is generally considered a hallmark of eryptosis, the degree of eryptosis was estimated based on the number of cells with low forward scatter (FSC) signal values. A comparison was made between the percentages of FSC-low erythrocytes [24].

Data analysis.

FlowJo™ (v10, BD Biosciences, USA) software was used to analyze data of eryptosis assays.

Statistical analysis.

ANOVA test was used to compare numerical data of eryptosis assays. Post-hoc Bonferroni test was applied afterwards. Data were processed by Graph Pad Prism 5.0 software (USA). The difference was considered statistically significant at $p < 0.05$.

The research was conducted in accordance with the fundamental principles of Good Clinical Practice (GCP, 1996), the Council of Europe Convention on Human Rights and Biomedicine (April 4, 1997), the Helsinki Declaration of the World Medical Association on Ethical Principles for Medical Research Involving Human Subjects (1964-2008), as well as the Ministry of Health of Ukraine Order No. 690 dated September 23, 2009 (amended by the Ministry of Health of Ukraine Order No. 523 dated July 12, 2012).

The research design was discussed and approved at the meeting of the Medical and Ethical Committee of the Kharkiv National Medical University (Protocol No. 19 dated May 19, 2024).

The paper is an excerpt from the scientific research project of the Department of Obstetrics and Gynecology No. 2 at Kharkiv National Medical University, entitled «Optimisation of diagnostics, treatment of diseases of the reproductive system and pregnancy complications in

women with extragenital pathology» (state registration number 0121U110923).

Results

The results of the study showed that during the examination of pregnant women met most common bacterial infections as *Escherichia coli* (45.5 %), *Staphylococcus epidermidis* (40.9 %), *Enterobacter aerogenes* (31.8 %), *Streptococcus agalactiae* (22.7 %), *Klebsiella pneumoniae* (13.6 %), *Ureaplasma urealiticum* (27.3 %), *Mycoplasma genitalium* (18.2 %); viral infections such as Cytomegalovirus (9.1 %), Herpes simplex virus (9.1 %), fungal infections *Candida albicans* – 59.1 %).

To determine whether genital tract infections affect the suicidal cell death modality of erythrocytes, key eryptotic parameters such as cellular morphology, asymmetry of phospholipid content, namely distribution of phosphatidylserine, in the plasma membrane of erythrocytes, and features of redox homeostasis were

evaluated. Detection was based on flow cytometry using commonly accepted biomarkers.

Importantly, phosphatidylserine externalization is considered a major biochemical sign of eryptosis that determines the clearance of circulating erythrocytes by phagocytic cells. In the current study, high Annexin V-FITC signaling indicated phosphatidylserine exposure and thus its location in the outer leaflet of the phospholipid bilayer. To quantify this process, we compared the percentage of cells displaying phosphatidylserine. The gating strategy is shown in Figure 1. Notably, fungal and viral reproductive tract infections during pregnancy are not associated with abnormal membrane phospholipid scrambling, as evidenced by the absence of statistically significant changes in the percentage of annexin V-positive cells (Figures 1 and 2a). Conversely, bacterial genital tract infections in pregnant women increased the percentage of circulating phosphatidylserine exposed cells, indicating activation of eryptosis (Figures 1 and 2a).




Fig. 1. Cell membrane phospholipid scrambling in circulating erythrocytes of pregnant women with genital tract infections. Representative histograms of pregnant women with bacterial (panel a), viral (panel b), fungal (panel c) reproductive tract infections and controls (panel d). The percentage of phosphatidylserine (PS)-exposing erythrocytes is statistically significantly higher in bacterial genital tract infections only. ANOVA and Bonferroni tests, Me and IQR (n = 10).

Note: * ($p < 0.05$) compared with the control group.

Oxidative stress is a common trigger for eryptosis. RBCs are known to be sensitive to ROS-mediated damage due to their inability to upregulate antioxidant enzymes. In this study, a ROS-sensitive H2DCDA probe was used to characterize the ROS-generating activity of red blood cells. Erythrocytes from pregnant patients with genital tract infections of bacterial, viral or fungal origin were found to

contain undetectable levels of intracellular ROS compared to control samples from healthy volunteers. As shown in Figures 2b and 3, the difference between the numerical values of the probe fluorescence, which characterizes the ROS content in erythrocytes, is negligible. Thus, oxidative stress is not observed in erythrocytes from patients with pregnancy-associated genital tract infections.




Fig. 2. Eryptosis-related parameters in pregnant women with bacterial (group 1), viral (group 2), fungal (group 3) reproductive tract infections and no pathology (group 4). Phosphatidylserine externalization (panel a), reactive oxygen species (ROS) production (panel b) and forward scatter (FSC) signaling (panel c) were quantified. ANOVA and Bonferroni tests, Me and IQR (n = 10). Note: * (p < 0.05) compared with the control group.




Fig. 3. Reactive oxygen species (ROS) production in red blood cells among pregnant females with bacterial (group 1), viral (group 2), fungal (group 3) reproductive tract infections and no genital tract infection (group 4). Representative histograms of ROS-sensitive probe's fluorescence demonstrate no changes in the redox homeostasis of circulating erythrocytes.

In addition to phospholipid bilayer asymmetry in cell membranes and oxidative stress, eryptosis is often associated with morphologic changes. Cell shrinkage is considered to be the most common abnormality in cell morphology observed in eryptosis. Flow cytometry-based detection of cell morphology focuses on the analysis of FSC signaling, which reflects cell shape. Our data show that reproductive tract infection in pregnant women is not associated with an increase in shrunken cells characterized by reduced FSC signaling (Figure 2c).

Discussion

In the current study, blood samples were collected from pregnant women with genital tract infections of bacterial, viral and fungal etiology to analyze key parameters of eryptosis. Adverse factors occurring in inflammatory conditions (circulating pro-inflammatory molecules or redox imbalance) are known to affect erythrocytes to induce their eryptosis, which may contribute to excessive erythrocyte loss, blood clotting, or abnormal rheological properties of blood [26]. However, eryptosis is generally considered to be protective because it occurs prior to hemolysis, preventing this pro-inflammatory accidental cell death mode of erythrocytes [23]. As has been widely reported, accelerated eryptosis may contribute to anemia in a variety of eryptosis-related diseases due to excessive erythrocyte clearance via externalized phosphatidylserine-

mediated efferocytosis [20, 27]. Indeed, our results clearly indicate that eryptosis may contribute to anemia in bacterial genital tract infections in pregnant women. Surprisingly, ROS-mediated pathways aren't involved in eryptosis associated with bacterial genital tract infections during pregnancy. However, bacterial urinary tract infections have been reported to be associated with impaired redox status [28]. Notably, oxidative stress is also observed at the systemic level in bacterial urinary tract infections in pregnant women and is manifested by elevation of circulating lipid peroxidation markers and depletion of the enzymatic linkage of the antioxidant system [29]. ROS overproduction has been suggested to be an important factor in the pathogenesis of pregnancy-associated urinary tract infections [30]. In eryptosis, ROS are known to mediate phosphatidylserine depletion by increasing scramblase activity and decreasing flippase activity, respectively [31]. However, this pathway was not activated in the present study. Thus, it is likely that other molecular mechanisms besides ROS are responsible for accelerated eryptosis in genital tract infections in pregnant women. In general, it is known that erythrocytes have a shorter lifespan and are more susceptible to destruction during pregnancy [32]. We believe that the pro-inflammatory environment associated with infections may facilitate erythrocyte elimination via eryptosis. Of note, viral and fungal infections have no effect on erythropoiesis

parameters in pregnant women, suggesting the pathogen-dependent nature of erythropoiesis stimulation.

Further exploration of the relationship between pregnancy-associated genital tract infections and eryptosis may provide a theoretical basis for targeting this cell death as a novel therapeutic approach to prevent anemia in pregnant women with this pathology.

Conclusions

This study demonstrates accelerated eryptosis in pregnancy-associated bacterial genital tract infections, which may provide a link between anemia and urinary tract infections. Eryptosis-

associated membrane phospholipid scrambling is observed exclusively in bacterial genital tract infections, whereas this hallmark of eryptosis is not seen in pregnant women with viral or fungal infections. Bacterial genital tract infection-induced eryptosis is ROS-independent and lacks cell shrinkage.

Conflict of Interests. The authors have no conflicts of interest to declare.

Founding source. This research received no specific grant from any funding agency in the public, commercial, or non-profit sectors.

References:

1. Marai W. Lower genital tract infections among pregnant women: a review. *East Afr Med J.* 2001;78(11):581-5.
2. Mobasheri M, Saeedi Varnamkhasht N, Karimi A, Banaeiyan S. Prevalence study of genital tract infections in pregnant women referred to health centers in Iran. *Turk J Med Sci.* 2014;44(2):232-6. doi: 10.3906/sag-1208-33
3. Zeng M, Yang L, Mao Y, He Y, Li M, Liu J, et al. Preconception reproductive tract infections status and adverse pregnancy outcomes: a population-based retrospective cohort study. *BMC Pregnancy and Childbirth [Internet].* 2022[cited 2024 Mar 10];22(1):501. Available from: <https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-022-04836-3> doi: 10.1186/s12884-022-04836-3
4. Zaytsev VT, Boyko VV, Parkhomenko KYu, Butkevich AYu. Sravnitel'naya otsenka techeniya peritonita, obuslovlennogo anaerobnymi neklostrioidal'nymi i aerobnymi mikroorganizmami [Comparative evaluation of the course of peritonitis caused by anaerobic non-clostridial and aerobic microorganisms]. *Klinichna khirurhiia.* 1999;1:5-6. (in Russian)
5. Laven J. Viral, bacterial and fungal infections in patients with recurrent miscarriage. *Human Reproduction [Internet].* 2023[cited 2024 Mar 10];38(S1): O-227. Available from: https://academic.oup.com/humrep/article/38/Supplement_1/dead093.277/7202557 doi: 10.1093/humrep/dead093.277
6. Liu J, Zeng M, Yang L, Mao Y, He Y, Li M, et al. Prevalence of reproductive tract infections among women preparing to conceive in Chongqing, China: trends and risk factors. *Reprod Health [Internet].* 2022[cited 2024 Mar 10];19(1):197. Available from: <https://reproductive-health-journal.biomedcentral.com/articles/10.1186/s12978-022-01502-x> doi: 10.1186/s12978-022-01502-x
7. Zeng M, Yang L, Mao Y, He Y, Li M, Liu J, et al. Preconception reproductive tract infections status and adverse pregnancy outcomes: a population-based retrospective cohort study. *BMC Pregnancy Childbirth [Internet].* 2022[cited 2024 Mar 10];22(1):501. Available from: <https://bmcpregnancychildbirth.biomedcentral.com/articles/10.1186/s12884-022-04836-3> doi: 10.1186/s12884-022-04836-3
8. Pasiëshvili NM, Vdovychenko YuP, Karpenko VH, Lazurenko VV. Patohenetychne obhruntuvannia systemnoi profilaktyky i terapii infektsiino-zapal'nykh urazhen' ploda zhinok, yaki strazhdaiut' na materyns'ko-plodovu infektsiiu [Pathogenetic substance of systemic prevention and therapy of infectious-inflammatory lesions of the fetus of women suffering maternal and fetal infection]. *Svit medytsyny ta biolohii.* 2019;2(68):96-100 doi: 10.26724/2079-8334-2019-2-68-96-100 (in Ukrainian).
9. Megli CJ, Coyne CB. Infections at the maternal-fetal interface: an overview of pathogenesis and defence. *Nat Rev Microbiol.* 2022;20(2):67-82. doi: 10.1038/s41579-021-00610-y
10. Stephen G, Mgongo M, Hussein Hashim T, Katanga J, Stray-Pedersen B, Msuya SE. Anaemia in Pregnancy: Prevalence, Risk Factors, and Adverse Perinatal Outcomes in Northern Tanzania. *Anemia [Internet].* 2018[cited 2024 Mar 10];2018:1846280. Available from: <https://downloads.hindawi.com/journals/anemia/2018/1846280.pdf> doi: 10.1155/2018/1846280
11. Pandey AK, Gautam D, Tolani H, Neogi SB. Clinical outcome post treatment of anemia in pregnancy with intravenous versus oral iron therapy: a systematic review and meta-analysis. *Sci Rep [Internet].* 2024[cited 2024 Mar 10];14(1):179. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10761955/pdf/41598_2023_Article_50234.pdf doi: 10.1038/s41598-023-50234-w
12. Tansarli GS, Karageorgopoulos DE, Kapaskelis A, Gkegkes I, Falagas ME. Iron deficiency and susceptibility to infections: evaluation of the clinical evidence. *Eur J Clin Microbiol Infect Dis.* 2013;32(10):1253-8. doi: 10.1007/s10096-013-1877-x
13. Dreischer P, Duszenko M, Stein J, Wieder T. Eryptosis: Programmed Death of Nucleus-Free, Iron-Filled Blood Cells. *Cells [Internet].* 2022[cited 2024 Mar 10];11(3):503. Available from: <https://www.mdpi.com/2073-4409/11/3/503> doi: 10.3390/cells11030503
14. Tkachenko A. Apoptosis and eryptosis: similarities and differences. *Apoptosis.* 2024;29(3-4):482-502. doi: 10.1007/s10495-023-01915-4
15. Tkachenko A, Onishchenko A. Casein kinase 1 α mediates eryptosis: a review. *Apoptosis.* 2023;28(1-2):1-19. doi: 10.1007/s10495-022-01776-3
16. Bissinger R, Qadri SM, Artunc F. Eryptosis: a driver of anemia in chronic kidney disease. *Curr Opin Nephrol Hypertens.* 2024;33(2):220-5. doi: 10.1097/mnh.0000000000000957
17. Alieksieieva O, Harkavenko K, Posokhov Y, Prokopyuk V, Lazurenko V, Safonov R. Redox status and cell membrane alterations of circulating leukocytes and erythrocytes in abnormal uterine bleeding. *Probl Endocrine Pathol.* 2023;80(1):7-16. doi: 10.21856/j-PEP.2023.1.01
18. Abdullaieva NA, Tertyshnyk DYu, Posokhov YeO, Lazurenko VV. Eryptosis in patients with gestational diabetes mellitus and hypertensive disorders of pregnancy. *Azerbaijan Medical Journal.* 2022;4:11-6. doi: 10.34921/amj.2022.4.022
19. Lang F, Gulbins E, Lang PA, Zappulla D, Föller M. Ceramide in suicidal death of erythrocytes. *Cell Physiol Biochem.* 2010;26(1):21-8. doi: 10.1159/000315102
20. Lang F, Qadri SM. Mechanisms and significance of eryptosis, the suicidal death of erythrocytes. *Blood Purif.* 2012;33(1-3):125-30. doi: 10.1159/000334163
21. Bissinger R, Bhuyan AAM, Qadri SM, Lang F. Oxidative stress, eryptosis and anemia: a pivotal mechanistic nexus in systemic diseases. *FEBS J.* 2019;286(5):826-54. doi: 10.1111/febs.14606

22. Kaliuzhka V, Tkachenko A, Myasoedov V, Markevych M, Onishchenko A, Babalyan I, et al. The Prognostic Value of Eryptosis Parameters in the Cerebrospinal Fluid for Cerebral Vasospasm and Delayed Cerebral Ischemia Formation. *World Neurosurg*[Internet]. 2023[cited 2024 Mar 5];173: e578-85. Available from: <https://www.sciencedirect.com/science/article/pii/S1878875023002449?via%3Dihub> doi: 10.1016/j.wneu.2023.02.096
23. Onishchenko AI, Prokopiuk VY, Chumachenko VA, Virych PA, Tryfonyuk LY, Kutsevol NV, et al. Hemocompatibility of dextran-graft-polyacrylamide/zinc oxide nanosystems: hemolysis or eryptosis? *Nanotechnology* [Internet]. 2023[cited 2024 Mar 10];35(3). Available from: <https://iopscience.iop.org/article/10.1088/1361-6528/ad02a3> doi: 10.1088/1361-6528/ad02a3
24. Yefimova S, Onishchenko A, Klochkov V, Myasoedov V, Kot Y, Tryfonyuk L, et al. Rare-earth orthovanadate nanoparticles trigger Ca²⁺-dependent eryptosis. *Nanotechnology* [Internet]. 2023[cited 2024 Mar 10];34(20). Available from: <https://iopscience.iop.org/article/10.1088/1361-6528/acbb7f> doi: 10.1088/1361-6528/acbb7f
25. Prokopiuk V, Yefimova S, Onishchenko A, Kapustnik V, Myasoedov V, Maksimchuk P, et al. Assessing the Cytotoxicity of TiO₂(2-x) Nanoparticles with a Different Ti(3+)(Ti(2+))/Ti(4+) Ratio. *Biol Trace Elem Res*. 2023;201(6):3117-30. doi: 10.1007/s12011-022-03403-3
26. Pretorius E. Erythrocyte deformability and eryptosis during inflammation, and impaired blood rheology. *Clin Hemorheol Microcirc*. 2018;69(4):545-50. doi: 10.3233/ch-189205
27. Lang F, Bissinger R, Abed M, Artunc F. Eryptosis – the Neglected Cause of Anemia in End Stage Renal Disease. *Kidney Blood Press Res*. 2017;42(4):749-60. doi: 10.1159/000484215
28. Chen Z, Zhang Z, Zhang H, Xie B. Analysis of the Oxidative Stress Status in Nonspecific Vaginitis and Its Role in Vaginal Epithelial Cells Apoptosis. *Biomed Res Int* [Internet]. 2015[cited 2024 Mar 10];2015:795656. Available from: <https://downloads.hindawi.com/journals/bmri/2015/795656.pdf> doi: 10.1155/2015/795656
29. Ciragil P, Kurutas EB, Gul M, Kilinc M, Aral M, Guven A. The effects of oxidative stress in urinary tract infection during pregnancy. *Mediators Inflamm*. 2005;2005(5):309-11. doi: 10.1155/mi.2005.309
30. Hussain T, Murtaza G, Metwally E, Kalhoro DH, Kalhoro MS, Rahu BA, et al. The Role of Oxidative Stress and Antioxidant Balance in Pregnancy. *Mediators Inflamm* [Internet]. 2021[cited 2024 Mar 10];2021:9962860. Available from: <https://downloads.hindawi.com/journals/mi/2021/9962860.pdf> doi: 10.1155/2021/9962860
31. Alghareeb SA, Alfihli MA, Fatima S. Molecular Mechanisms and Pathophysiological Significance of Eryptosis. *Int J Mol Sci* [Internet]. 2023[cited 2024 Mar 10];24(6). Available from: <https://www.mdpi.com/1422-0067/24/6/5079> doi: 10.3390/ijms24065079
32. Lurie S, Mamet Y. Red blood cell survival and kinetics during pregnancy. *European J Obstet Gynecol Reprod Biol*. 2000;93(2):185-92. doi: 10.1016/S0301-2115(00)00290-6

ЕРИПТОЗ ЯК ЗВ'ЯЗОК МІЖ БАКТЕРІАЛЬНОЮ ІНФЕКЦІЄЮ РЕПРОДУКТИВНОГО ТРАКТА ТА АНЕМІЄЮ ВАГІТНИХ

І. Д. Кудін, В. В. Лазуренко, В. Ю. Прокопюк, О. Ю. Железняков, М. Л. Панасовський

**Харківський національний медичний університет.
(м. Харків, Україна)**

Резюме.

Інфекції генітального тракту різної етіології і анемія залишаються поширеними серед вагітних жінок. На даний момент існують мізерні дані про вплив інфекцій репродуктивного тракту на анемію, яка розвивається у жінок під час вагітності, особливо фокусуючись на процесах ериптозу.

Метою даного дослідження було оцінити здатність бактеріальної, вірусної та грибової інфекції репродуктивного тракту стати тригером процесів ериптозу циркулюючих еритроцитів у вагітних жінок.

Матеріали і методи. Було обстежено 47 вагітних, яких в залежності від наявності інфекційної патології було поділено на три групи: 14 вагітних з бактеріальним ураженням, 12 жінок з вірусним ураженням, 11 вагітних з грибовою інфекцією, 10 жінок без ознак інфікування склали контрольну групу. Для визначення інфікування використовували наступні методи діагностики: бактеріологічний, мікроскопічний, імуоферментний, полімеразну ланцюгову реакцію. За допомогою проточної цитометрії визначали параметри ериптозу (клітинну морфологію, скремблінг фосфоліпідної мембрани та оксидативний стрес) в крові вагітних з бактеріальною, вірусною, грибовою інфекцією репродуктивного тракту. Фарбування анексин V-FITC та 2',7'-діхлориддигідрофлуоресцина діацетата (H2DCFDA) використовувалося для кількісного визначення фосфатиділсерину і утворення активних форм кисню (АФК) відповідно Флуоресценція визначалася за допомогою наборів BD FACSCanto™ II для проточної цитометрії.

Тест ANOVA використовувався для порівняння числових даних аналізів ериптозу. Після цього був застосований тест Бонферроні. Дані обробляли програмою Graph Pad Prism 5.0 (США). Різницю вважали статистично значущою при $p < 0,05$.

Дослідження проводилися відповідно до основоположних принципів належної клінічної практики (GCP, 1996), Конвенції Ради Європи про права людини та біомедицину (4 квітня 1997 року), Гельсінської декларації Всесвітньої медичної асоціації про етичні принципи медичних Дослідження на людях (1964-2008), а також наказ МОЗ України від 23.09.2009 № 690 (зміни внесені наказом МОЗ України від 12.07.2012 № 523). Проект дослідження обговорено та схвалено на засіданні медико-етичної комісії ХНМУ (протокол № 19 від 19 травня 2024 р.).

Стаття є фрагментом наукового проекту кафедри акушерства та гінекології № 2 Харківського національного медичного університету «Оптимізація діагностики, лікування захворювань репродуктивної системи та ускладнень вагітності у жінок з екстрагенітальною патологією» (державний реєстраційний номер 0121U110923).

Результати. Параметри ериптозу були оцінені у вагітних жінок з бактеріальним, вірусним, грибовим інфікуванням репродуктивного тракту. Ураження репродуктивного тракту вагітних бактеріальною флорою супроводжувалося підсиленням екстерналізації фосфатидилсерину серед циркулюючих еритроцитів з/без накопичуванням АФК та зморщуванням клітин. Еритроцити від вагітних з вірусним або грибовим ураженням не продемонстрували ознак ериптозу, таких як скремблінг клітинних мембран, оксидативний стрес або зміни клітинної морфології.

Висновки. Інфекції репродуктивного тракту бактеріального походження у вагітних жінок пов'язані з прискоренням процесів ериптозу, який може призводити до анемії у цих вагітних. Виключно при бактеріальній інфекції генітального тракту було показано, що запускається скремблуння і з цього часу ериптоз в еритроцитах вагітних жінок. Екстерналізація фосфатиділсерину незначно відрізнялися серед вірусної або грибової інфекцій та здоровими вагітними групи контролю. Бактеріальне інфікування генітального тракту індукує процеси ериптозу, які не були пов'язані ні з усадкою клітин, ні з надлишковою продукцією АФК.

Ключові слова: інфекції репродуктивного тракту; еритроцити; анемія; вагітність; ериптоз.

Contact Information:

Illia Kudin – Postgraduate student of the Department of Obstetrics and Gynecology № 2, Kharkiv National Medical University (Kharkiv, Ukraine)

e-mail: idkudin.po23@knmu.edu.ua

ORCID ID: <https://orcid.org/0009-0001-7159-4796>

Viktoriya Lazurenko – Doctor of Medical Sciences, Professor, Head of the Department of Obstetrics and Gynecology № 2, Kharkiv National Medical University (Kharkiv, Ukraine)

e-mail: vv.lazurenko@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0002-7300-4868>

Researcher ID: https://www.researchgate.net/profile/Viktoriya_Lazurenko/experience

Scopus Author ID: <https://www.scopus.com/%20authrid/detail.uri?authorid=57194712400>

Volodymyr Prokopiuk – Candidate of Medical Sciences, Director of the Research Institute of Experimental and Clinical Medicine, Kharkiv National Medical University (Kharkiv, Ukraine)

e-mail: vy.prokopiuk@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0003-4379-4130>

Researcher ID: <https://www.researchgate.net/profile/Volodymyr-Prokopyuk>

Scopus Author ID: <https://www.scopus.com/authid/detail.uri?authorid=56964479700>

Oleksandr Zhelezniakov – Candidate of Medical Sciences, Associate Professor of the Department of Obstetrics and Gynecology № 3, Kharkiv National Medical University (Kharkiv, Ukraine)

e-mail: oy.zhelezniakov@knmu.edu.ua

ORCID ID: <https://orcid.org/0009-0004-4667-9191>

Mykola Panasovskyi – Doctor of Medical Sciences, Professor of the Department of Department of Urology, Nephrology and Andrology named after prof. A. G. Podreza, Kharkiv National Medical University (Kharkiv, Ukraine)

e-mail: ml.panasovskyi@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0001-7152-0204>

Scopus Author ID: <https://www.scopus.com/authid/detail.uri?authorid=57219602673>

Контактна інформація:

Кудін Ілля Дмитрович – аспірант кафедри акушерства та гінекології № 2, Харківський національний медичний університет (Харків, Україна)

e-mail: idkudin.po23@knmu.edu.ua

ORCID ID: <https://orcid.org/0009-0001-7159-4796>

Лазуренко Вікторія Валентинівна – доктор медичних наук, професор, завідувачка кафедри акушерства та гінекології № 2, Харківський національний медичний університет (Харків, Україна)

e-mail: vv.lazurenko@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0002-7300-4868>

Researcher ID: https://www.researchgate.net/profile/Viktoriya_Lazurenko/experience

Scopus Author ID: <https://www.scopus.com/%20authrid/detail.uri?authorid=57194712400>

Прокопюк Володимир Юрійович – кандидат медичних наук, директор науково-дослідного інституту експериментальної та клінічної медицини, Харківський національний медичний університет (Харків, Україна)

e-mail: vy.prokopiuk@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0003-4379-4130>

Researcher ID: <https://www.researchgate.net/profile/Volodymyr-Prokopyuk>

Scopus Author ID: <https://www.scopus.com/authid/detail.uri?authorid=56964479700>

Железняков Олександр Юрійович – кандидат медичних наук, доцент кафедри акушерства та гінекології № 3, Харківський національний медичний університет (Харків, Україна)

e-mail: oy.zhelezniakov@knmu.edu.ua

ORCID ID: <https://orcid.org/0009-0004-4667-9191>

Панасовський Микола Леонідович – доктор медичних наук, професор кафедри урології, нефрології та андрології імені проф. А. Г. Подреза, Харківський національний медичний університет (Харків, Україна)

e-mail: ml.panasovskyi@knmu.edu.ua

ORCID ID: <https://orcid.org/0000-0001-7152-0204>

Scopus Author ID: <https://www.scopus.com/authid/detail.uri?authorid=57219602673>



Received for editorial office on 10/02/2024

Signed for printing on 14/04/2024